

Rabbit CYB5R3 Polyclonal Antibody | anti-CYB5R3 antibody

Anti-CYB5R3 Antibody Picoband

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA126312	Rabbit	Human, Mouse, Rat	ELISA, Flow Cytometry, Functional Assay, Immunofluorescence, Immunocytochemistry, Western Blot


PRODUCT SYNONYMS

CYB5R3; Polyclonal Antibody; Anti-CYB5R3 Antibody Picoband; anti-CYB5R3 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	Rabbit IgG
Purity/Purification	Immunogen affinity purified.
Form/Format	Lyophilized Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml. (varies by lot)
Applicable Applications for anti-CYB5R3 antibody	ELISA, FCM/FACS (Flow Cytometry), IF (Immunofluorescence), ICC (Immunocytochemistry), WB (Western Blot)
Reconstitution	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Immunogen	E Coli-derived human CYB5R3 recombinant protein (Position: M24-F301).
Preparation and Storage	Store at -20 degree C for one year from date of receipt. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for six months. Avoid repeated freezing and thawing.

IMAGES



FCM/FACS (Flow Cytometry)

Figure 4. Flow Cytometry analysis of 293T cells using anti-CYB5R3 antibody (AAA126312). Overlay histogram showing 293T cells stained with AAA126312 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-CYB5R3 Antibody (AAA126312, 1 ug/1x10⁶ cells) for 30 min at 20 degree C. DyLight488 conjugated goat anti-rabbit IgG was used as secondary antibody for 30 minutes at 20 degree C. Isotype control antibody (Green line) was rabbit IgG (1 ug/1x10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.



IF (Immunofluorescence)

Figure 3. IF analysis of CYB5R3 using anti-CYB5R3 antibody (AAA126312). CYB5R3 was detected in an immunocytochemical section of HEP3B cells. Enzyme antigen retrieval was performed using IHC enzyme antigen retrieval reagent (AR0022) for 15 mins. The cells were blocked with 10% goat serum. And then incubated with 5 ug/mL rabbit anti-CYB5R3 Antibody (AAA126312) overnight at 4 degree C. DyLight488 Conjugated Goat Anti-Rabbit IgG was used as secondary antibody at 1:100 dilution and incubated for 30 minutes at 37 degree C. The section was counterstained with DAPI. Visualize using a fluorescence microscope and filter sets appropriate for the label used.



IF (Immunofluorescence)

Figure 2. IF analysis of CYB5R3 using anti-CYB5R3 antibody (AAA126312).CYB5R3 was detected in an immunocytochemical section of U2OS cells. Enzyme antigen retrieval was performed using IHC enzyme antigen retrieval reagent (AR0022) for 15 mins. The cells were blocked with 10% goat serum. And then incubated with 5 ug/mL rabbit anti-CYB5R3 Antibody (AAA126312) overnight at 4 degree C. DyLight488 Conjugated Goat Anti-Rabbit IgG was used as secondary antibody at 1:100 dilution and incubated for 30 minutes at 37 degree C. The section was counterstained with DAPI. Visualize using a fluorescence microscope and filter sets appropriate for the label used.



WB (Western Blot)

Figure 1. Western blot analysis of CYB5R3 using anti-CYB5R3 antibody (AAA126312).Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel)/90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.Lane 1: human HepG2 whole cell lysates,Lane 2: human A431 whole cell lysates,Lane 3: rat liver tissue lysates,Lane 4: rat lung tissue lysates,Lane 5: mouse liver tissue lysates,Lane 6: mouse lung tissue lysates.After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-CYB5R3 antigen affinity purified polyclonal antibody (#AAA126312) at 0.25 ug/mL overnight at 4 degree C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for CYB5R3 at approximately 34 kDa. The expected band size for CYB5R3 is at 34 kDa.

ADDITIONAL INFORMATION

Related Product Information for anti-CYB5R3 antibody	NADH-cytochrome b5 reductase 3 is an enzyme that in humans is encoded by the CYB5R3 gene. This gene encodes cytochrome b5 reductase, which includes a membrane-bound form in somatic cells (anchored in the endoplasmic reticulum, mitochondrial and other membranes) and a soluble form in erythrocytes. The membrane-bound form exists mainly on the cytoplasmic side of the endoplasmic reticulum and functions in desaturation and elongation of fatty acids, in cholesterol biosynthesis, and in drug metabolism. The erythrocyte form is located in a soluble fraction of circulating erythrocytes and is involved in methemoglobin reduction. The membrane-bound form has both membrane-binding and catalytic domains, while the soluble form has only the catalytic domain. Alternate splicing results in multiple transcript variants. Mutations in this gene cause methemoglobinemias.
Product Categories/Family for anti-CYB5R3 antibody	Primary Antibodies, Rabbit Polyclonal Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	4503327
NCBI GeneID	1727
NCBI Accession #	NP_000389.1
NCBI GenBank Nucleotide #	NM_000398.6
UniProt Accession #	P00387; O75675; Q8TDL8; Q8WTS8; Q9UEN4; Q9UEN5; Q9UL55; Q9UL56; B1AHF2; B7Z7L3
Molecular Weight	34,235 Da
NCBI Official Full Name	NADH-cytochrome b5 reductase 3 isoform 1
NCBI Official Synonym Full Names	cytochrome b5 reductase 3
NCBI Official Symbol	CYB5R3
NCBI Official Synonym Symbols	B5R; DIA1
NCBI Protein Information	NADH-cytochrome b5 reductase 3; diaphorase-1; NADH-cytochrome b5 reductase 3 soluble form; NADH-cytochrome b5 reductase 3 membrane-bound form
UniProt Protein Name	NADH-cytochrome b5 reductase 3
UniProt Gene Name	CYB5R3
UniProt Synonym Gene Names	DIA1; B5R
UniProt Entry Name	NB5R3_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/cyb5r3-antibody-flow-cytometry-immunofluorescence-immunocytochemistry-western-blot-elisa/126312>