

Rabbit CPT1B Polyclonal Antibody | anti-CPT1B antibody

Anti-CPT1B Antibody Picoband


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA126318	Rabbit	Human, Mouse, Rat, Monkey	ELISA, Flow Cytometry, Functional Assay, Immunofluorescence, Immunocytochemistry, Western Blot

PRODUCT SYNONYMS


CPT1B; Polyclonal Antibody; Anti-CPT1B Antibody Picoband; anti-CPT1B antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat, Monkey
Clonality	Polyclonal
Isotype	Rabbit IgG
Purity/Purification	Immunogen affinity purified.
Form/Format	Lyophilized Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml. (varies by lot)
Applicable Applications for anti-CPT1B antibody	ELISA, FCM/FACS (Flow Cytometry), IF (Immunofluorescence), ICC (Immunocytochemistry), WB (Western Blot)
Reconstitution	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Immunogen	E Coli-derived human CPT1B recombinant protein (Position: R180-S772).
Preparation and Storage	Store at -20 degree C for one year from date of receipt. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for six months. Avoid repeated freezing and thawing.



FCM/FACS (Flow Cytometry)
Figure 3. Flow Cytometry analysis of HeLa cells using anti-CPT1B antibody (AAA126318).Overlay histogram showing HeLa cells stained with AAA126318 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-CPT1B Antibody (AAA126318, 1 ug/1x10⁶ cells) for 30 min at 20 degree C. DyLight488 conjugated goat anti-rabbit IgG was used as secondary antibody for 30 minutes at 20 degree C. Isotype control antibody (Green line) was rabbit IgG (1 ug/1x10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.



IF (Immunofluorescence)
Figure 2. IF analysis of CPT1B using anti-CPT1B antibody (AAA126318).CPT1B was detected in an immunocytochemical section of HeLa cells. Enzyme antigen retrieval was performed using IHC enzyme antigen retrieval reagent (AR0022) for 15 mins. The cells were blocked with 10% goat serum. And then incubated with 5 ug/mL rabbit anti-CPT1B Antibody (AAA126318) overnight at 4 degree C. DyLight488 Conjugated Goat Anti-Rabbit IgG was used as secondary antibody at 1:100 dilution and incubated for 30 minutes at 37 degree C. The section was counterstained with DAPI. Visualize using a fluorescence microscope and filter sets appropriate for the label used.



WB (Western Blot)
Figure 1. Western blot analysis of CPT1B using anti-CPT1B antibody (AAA126318).Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel)/90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.Lane 1: monkey heart tissue lysates,Lane 2: rat heart tissue lysates,Lane 3: mouse heart tissue lysates.After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-CPT1B antigen affinity purified polyclonal antibody (#AAA126318) at 0.5 ug/mL overnight at 4 degree C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for CPT1B at approximately 88 kDa. The expected band size for CPT1B is at 88 kDa.

ADDITIONAL INFORMATION

Related Product Information for anti-CPT1B antibody	CPT1B is located on 22q13.33. The protein encoded by this gene, a member of the carnitine/ choline acetyltransferase family, is the rate-controlling enzyme of the long-chain fatty acid beta-oxidation pathway in muscle mitochondria. This enzyme is required for the net transport of long-chain fatty acyl-CoAs from the cytoplasm into the mitochondria. Multiple transcript variants encoding different isoforms have been found for this gene, and read-through transcripts are expressed from the upstream locus that include exons from this gene.
Product Categories/Family for anti-CPT1B antibody	Primary Antibodies, Rabbit Polyclonal Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	223468670
NCBI GeneID	1375
NCBI Accession #	NP_001138606.1
NCBI GenBank Nucleotide #	NM_001145134.1
UniProt Accession #	Q92523; Q13389; Q99655; Q9BY90; B7Z4U4; B7Z5T8; E9PCP2
Molecular Weight	87,801 Da
NCBI Official Full Name	carnitine O-palmitoyltransferase 1, muscle isoform isoform c
NCBI Official Synonym Full Names	carnitine palmitoyltransferase 1B (muscle)
NCBI Official Symbol	CPT1B
NCBI Official Synonym Symbols	CPTI; CPT1M; MCPT1; CPT1-M; CPTI-M; M-CPT1; MCCPT1
NCBI Protein Information	carnitine O-palmitoyltransferase 1, muscle isoform; carnitine palmitoyltransferase I-like protein; carnitine O-palmitoyltransferase I, mitochondrial muscle isoform
UniProt Protein Name	Carnitine O-palmitoyltransferase 1, muscle isoform
UniProt Gene Name	CPT1B
UniProt Synonym Gene Names	KIAA1670; CPT1-M; CPT I; CPTI-M
UniProt Entry Name	CPT1B_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/cpt1b-antibody-flow-cytometry-immunofluorescence-immunocytochemistry-western-blot-elisa/126318>