

Rabbit anti-Human ARL1 Polyclonal Antibody | anti-ARL1 antibody

Anti-ARL1 Antibody Picoband

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA126333	Rabbit	Human	Flow Cytometry, Functional Assay, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

ARL1; Polyclonal Antibody; Anti-ARL1 Antibody Picoband; anti-ARL1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Purity/Purification	Immunogen affinity purified.
Form/Format	Lyophilized Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml. (varies by lot)
Applicable Applications for anti-ARL1 antibody	FCM/FACS (Flow Cytometry), IHC (Immunohistochemistry), WB (Western Blot)
Reconstitution	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Immunogen	A synthetic peptide corresponding to a sequence of human ARL1 (LDEAMEWLVELKSRQ).
Preparation and Storage	Store at -20 degree C for one year from date of receipt. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for six months. Avoid repeated freezing and thawing.

IMAGES




FCM/FACS (Flow Cytometry)

Figure 4. Flow Cytometry analysis of RT4 cells using anti-ARL1 antibody (AAA126333). Overlay histogram showing RT4 cells stained with AAA126333 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-ARL1 Antibody (AAA126333, 1 ug/1x10⁶ cells) for 30 min at 20 degree C. DyLight488 conjugated goat anti-rabbit IgG was used as secondary antibody for 30 minutes at 20 degree C. Isotype control antibody (Green line) was rabbit IgG (1 ug/1x10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.




IHC (Immunohistochemistry)

Figure 3. IHC analysis of ARL1 using anti-ARL1 antibody (AAA126333). ARL1 was detected in a paraffin-embedded section of human colorectal adenocarcinoma tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-ARL1 Antibody (AAA126333) overnight at 4 degree C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit with DAB as the chromogen.



IHC (Immunohistochemistry)

Figure 2. IHC analysis of ARL1 using anti-ARL1 antibody (AAA126333). ARL1 was detected in a paraffin-embedded section of human cervical cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-ARL1 Antibody (AAA126333) overnight at 4 degree C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit with DAB as the chromogen.



WB (Western Blot)

Figure 1. Western blot analysis of ARL1 using anti-ARL1 antibody (AAA126333). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel)/90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: human K562 whole cell lysates, Lane 2: human Hep2 whole cell lysates, Lane 3: human HepG2 whole cell lysates, Lane 4: human RT4 whole cell lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-ARL1 antigen affinity purified polyclonal antibody (#AAA126333) at 0.5 ug/mL overnight at 4 degree C, then washed with TBS-0.1% Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for ARL1 at approximately 20 kDa. The expected band size for ARL1 is at 20 kDa.

ADDITIONAL INFORMATION

Related Product Information for anti-ARL1 antibody	ADP-ribosylation factor-like protein 1 is a protein that in humans is encoded by the ARL1 gene. The protein encoded by this gene belongs to the ARL (ADP-ribosylation factor-like) family of proteins, which are structurally related to ADP-ribosylation factors (ARFs). ARFs, described as activators of cholera toxin (CT) ADP-ribosyltransferase activity, regulate intracellular vesicular membrane trafficking, and stimulate a phospholipase D (PLD) isoform. Although, ARL proteins were initially thought not to activate CT or PLD, later work showed that they are weak stimulators of PLD and CT in a phospholipid dependent manner. Alternative splicing results in multiple transcript variants encoding different isoforms.
Product Categories/Family for anti-ARL1 antibody	Primary Antibodies , Rabbit Polyclonal Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	728888
NCBI GeneID	400
UniProt Accession #	P40616 ; P80417 ; Q53XB1 ; B4DWW1
Molecular Weight	181
NCBI Official Full Name	ADP-ribosylation factor-like protein 1
NCBI Official Synonym Full Names	ADP-ribosylation factor-like 1
NCBI Official Symbol	ARL1
NCBI Official Synonym Symbols	ARFL1
NCBI Protein Information	ADP-ribosylation factor-like protein 1
UniProt Protein Name	ADP-ribosylation factor-like protein 1
UniProt Gene Name	ARL1

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/ar1-antibody-flow-cytometry-immunohistochemistry-western-blot-elisa/126333>