

Rabbit MEK5/MAP2K5 Polyclonal Antibody | anti-MAP2K5 antibody

Anti-MEK5/MAP2K5 Antibody Picoband

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA126346	Rabbit	Human, Mouse, Rat	ELISA, Immunofluorescence, Immunocytochemistry, Western Blot

PRODUCT SYNONYMS

MEK5/MAP2K5; Polyclonal Antibody; Anti-MEK5/MAP2K5 Antibody Picoband; anti-MAP2K5 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	Rabbit IgG
Purity/Purification	Immunogen affinity purified.
Form/Format	Lyophilized Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml. (varies by lot)
Applicable Applications for anti-MAP2K5 antibody	ELISA, IF (Immunofluorescence), ICC (Immunocytochemistry), WB (Western Blot)
Reconstitution	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Immunogen	E Coli-derived human MEK5/MAP2K5 recombinant protein (Position: R401-Q445).
Preparation and Storage	Store at -20 degree C for one year from date of receipt. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for six months. Avoid repeated freezing and thawing.

IMAGES



IF (Immunofluorescence)

Figure 2. IF analysis of MEK5/MAP2K5 using anti-MEK5/MAP2K5 antibody (AAA126346). MEK5/MAP2K5 was detected in an immunocytochemical section of A549 cells. Enzyme antigen retrieval was performed using IHC enzyme antigen retrieval reagent (AR0022) for 15 mins. The cells were blocked with 10% goat serum. And then incubated with 5 ug/mL rabbit anti-MEK5/MAP2K5 Antibody (AAA126346) overnight at 4 degree C. DyLight594 Conjugated Goat Anti-Rabbit IgG (BA1142) was used as secondary antibody at 1:100 dilution and incubated for 30 minutes at 37 degree C. The section was counterstained with DAPI. Visualize using a fluorescence microscope and filter sets appropriate for the label used.



WB (Western Blot)

Figure 1. Western blot analysis of MEK5/MAP2K5 using anti-MEK5/MAP2K5 antibody (AAA126346). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel)/90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: human Jurkat whole cell lysates, Lane 2: rat heart tissue lysates, Lane 3: rat skeletal muscle tissue lysates, Lane 4: mouse heart tissue lysates, Lane 5: mouse skeletal muscle tissue lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-MEK5/MAP2K5 antigen affinity purified polyclonal antibody (#AAA126346) at 0.5 ug/mL overnight at 4 degree C, then washed with TBS-0.1% Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for MEK5/MAP2K5 at approximately 50 kDa. The expected band size for MEK5/MAP2K5 is at 50 kDa.

ADDITIONAL INFORMATION

Related Product Information for anti-MAP2K5 antibody	Dual specificity mitogen-activated protein kinase kinase 5 is an enzyme that in humans is encoded by the MAP2K5 gene. The protein encoded by this gene is a dual specificity protein kinase that belongs to the MAP kinase kinase family. This kinase specifically interacts with and activates MAPK7/ERK5. This kinase itself can be phosphorylated and activated by MAP3K3/MEKK3, as well as by atypical protein kinase C isoforms (aPKCs). The signal cascade mediated by this kinase is involved in growth factor stimulated cell proliferation and muscle cell differentiation. Three alternatively spliced transcript variants of this gene encoding distinct isoforms have been described.
Product Categories/Family for anti-MAP2K5 antibody	Primary Antibodies, Rabbit Polyclonal Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	332164798
NCBI GeneID	5607
NCBI Accession #	NP_001193733.1
NCBI GenBank Nucleotide #	NM_001206804.1
UniProt Accession #	Q13163; Q92961; Q92962; B4DE43
Molecular Weight	50112 MW
NCBI Official Full Name	dual specificity mitogen-activated protein kinase kinase 5 isoform C
NCBI Official Synonym Full Names	mitogen-activated protein kinase kinase 5
NCBI Official Symbol	MAP2K5
NCBI Official Synonym Symbols	MEK5; MAPKK5; PRKMK5; HsT17454
NCBI Protein Information	dual specificity mitogen-activated protein kinase kinase 5
UniProt Protein Name	Dual specificity mitogen-activated protein kinase kinase 5
UniProt Gene Name	MAP2K5
UniProt Synonym Gene Names	MEK5; MKK5; PRKMK5; MAP kinase kinase 5; MAPKK 5; MEK 5

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/mek5-map2k5-antibody-immunofluorescence-immunocytochemistry-western-blot-elisa/126346>