

Rabbit Azurocidin/AZU1 Polyclonal Antibody | anti-AZU1 antibody

Anti-Azurocidin/AZU1 Antibody Picoband

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA126416	Rabbit	Human, Mouse, Rat	ELISA, Flow Cytometry, Functional Assay, Immunohistochemistry, Western Blot

PRODUCT SYNONYMS

Azurocidin/AZU1; Polyclonal Antibody; Anti-Azurocidin/AZU1 Antibody Picoband; anti-AZU1 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	Rabbit IgG
Purity/Purification	Immunogen affinity purified.
Form/Format	Lyophilized Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml. (varies by lot)
Applicable Applications for anti-AZU1 antibody	ELISA, FCM/FACS (Flow Cytometry), IHC (Immunohistochemistry), WB (Western Blot)
Reconstitution	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Immunogen	E Coli-derived human Azurocidin/AZU1 recombinant protein (Position: I27-N245).
Preparation and Storage	Store at -20 degree C for one year from date of receipt. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for six months. Avoid repeated freezing and thawing.

IMAGES



FCM/FACS (Flow Cytometry)

Figure 3. Flow Cytometry analysis of THP-1 cells using anti-Azurocidin/AZU1 antibody (AAA126416). Overlay histogram showing THP-1 cells stained with AAA126416 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-Azurocidin/AZU1 Antibody (AAA126416, 1 ug/1x10⁶ cells) for 30 min at 20 degree C. DyLight488 conjugated goat anti-rabbit IgG was used as secondary antibody for 30 minutes at 20 degree C. Isotype control antibody (Green line) was rabbit IgG (1 ug/1x10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.



IHC (Immunohistochemistry)

Figure 2. IHC analysis of Azurocidin/AZU1 using anti-Azurocidin/AZU1 antibody (AAA126416). Azurocidin/AZU1 was detected in a paraffin-embedded section of human appendicitis tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-Azurocidin/AZU1 Antibody (AAA126416) overnight at 4 degree C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit with DAB as the chromogen.



WB (Western Blot)

Figure 1. Western blot analysis of Azurocidin/AZU1 using anti-Azurocidin/AZU1 antibody (AAA126416). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel)/90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: rat spleen tissue lysates, Lane 2: rat thymus tissue lysates, Lane 3: mouse spleen tissue lysates, Lane 4: mouse thymus tissue lysates, Lane 5: mouse RAW264.7 whole cell lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-Azurocidin/AZU1 antigen affinity purified polyclonal antibody (#AAA126416) at 0.5 ug/mL overnight at 4 degree C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for Azurocidin/AZU1 at approximately 37 kDa. The expected band size for Azurocidin/AZU1 is at 37 kDa.

ADDITIONAL INFORMATION

Related Product Information for anti-AZU1 antibody	Azurocidin also known as cationic antimicrobial protein CAP37 or heparin-binding protein (HBP) is a protein that in humans is encoded by the AZU1 gene. Azurophil granules, specialized lysosomes of the neutrophil, contain at least 10 proteins implicated in the killing of microorganisms. This gene encodes a preproprotein that is proteolytically processed to generate a mature azurophil granule antibiotic protein, with monocyte chemotactic and antimicrobial activity. It is also an important multifunctional inflammatory mediator. This encoded protein is a member of the serine protease gene family but it is not a serine proteinase, because the active site serine and histidine residues are replaced. The genes encoding this protein, neutrophil elastase 2, and proteinase 3 are in a cluster located at chromosome 19pter. All 3 genes are expressed coordinately and their protein products are packaged together into azurophil granules during neutrophil differentiation.
Product Categories/Family for anti-AZU1 antibody	Primary Antibodies, Rabbit Polyclonal Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	11342670
NCBI GeneID	566
NCBI Accession #	NP_001691.1
NCBI GenBank Nucleotide #	NM_001700.3
UniProt Accession #	P20160; P80014; Q52LG4; Q9UCM1; Q9UCT5
Molecular Weight	26,886 Da
NCBI Official Full Name	azurocidin preproprotein
NCBI Official Synonym Full Names	azurocidin 1
NCBI Official Symbol	AZU1
NCBI Official Synonym Symbols	AZU; HBP; NAZC; AZAMP; CAP37; HUMAZUR
NCBI Protein Information	azurocidin; neutrophil azurocidin; heparin-binding protein; cationic antimicrobial protein 37; cationic antimicrobial protein CAP37
UniProt Protein Name	Azurocidin
UniProt Gene Name	AZU1
UniProt Synonym Gene Names	HBP
UniProt Entry Name	CAP7_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/azurocidin-azu1-antibody-flow-cytometry-immunohistochemistry-western-blot-elisa/126416>