

Rabbit anti-Human Calponin 2/CNN2 Polyclonal Antibody | anti-CNN2 antibody

Anti-Calponin 2/CNN2 Antibody Picoband

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA126530	Rabbit	Human	Western Blot, Immunohistochemistry, Flow Cytometry, Functional Assay


PRODUCT SYNONYMS

Calponin 2/CNN2; Polyclonal Antibody; Anti-Calponin 2/CNN2 Antibody Picoband; anti-CNN2 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Purity/Purification	Immunogen affinity purified.
Form/Format	Lyophilized Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml. (varies by lot)
Applicable Applications for anti-CNN2 antibody	WB (Western Blot), IHC (Immunohistochemistry), FCM/FACS (Flow Cytometry)
Reconstitution	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Immunogen	A synthetic peptide corresponding to a sequence of human Calponin 2/CNN2 (DCPDGPGEVPEYPPYYQEEAGY).
Preparation and Storage	Store at -20 degree C for one year from date of receipt. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for six months. Avoid repeated freezing and thawing.

IMAGES




FCM/FACS (Flow Cytometry)

Figure 3. Flow Cytometry analysis of Daudi cells using anti-Calponin 2/CNN2 antibody (AAA126530). Overlay histogram showing Daudi cells stained with AAA126530 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-Calponin 2/CNN2 Antibody (AAA126530, 1 ug/1x10⁶ cells) for 30 min at 20 degree C. DyLight488 conjugated goat anti-rabbit IgG was used as secondary antibody for 30 minutes at 20 degree C. Isotype control antibody (Green line) was rabbit IgG (1 ug/1x10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.



IHC (Immunohistochemistry)

Figure 2. IHC analysis of Calponin 2/CNN2 using anti-Calponin 2/CNN2 antibody (AAA126530). Calponin 2/CNN2 was detected in a paraffin-embedded section of human tonsil tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-Calponin 2/CNN2 Antibody (AAA126530) overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.



WB (Western Blot)

Figure 1. Western blot analysis of Calponin 2/CNN2 using anti-Calponin 2/CNN2 antibody (AAA126530). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel)/90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: human HepG2 whole cell lysates, Lane 2: human HeLa whole cell lysates, Lane 3: human RT4 whole cell lysates, Lane 4: human Caco-2 whole cell lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-Calponin 2/CNN2 antigen affinity purified polyclonal antibody (#AAA126530) at 0.25 ug/mL overnight at 4 degree C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for Calponin 2/CNN2 at approximately 34 kDa. The expected band size for Calponin 2/CNN2 is at 34 kDa.

ADDITIONAL INFORMATION

Related Product Information for anti-CNN2 antibody	Calponin 2 is a protein that in humans is encoded by the CNN2 gene. The protein encoded by this gene, which can bind actin, calmodulin, troponin C, and tropomyosin, may function in the structural organization of actin filaments. The encoded protein could play a role in smooth muscle contraction and cell adhesion. Several pseudogenes of this gene have been identified, and are present on chromosomes 1, 2, 3, 6, 9, 11, 13, 15, 16, 21 and 22. Alternative splicing results in multiple transcript variants encoding different isoforms.
Product Categories/Family for anti-CNN2 antibody	Primary Antibodies , Rabbit Polyclonal Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	4758018
NCBI GeneID	1265
NCBI Accession #	NP_004359.1
NCBI GenBank Nucleotide #	NM_004368.2
UniProt Accession #	Q99439 ; Q92578 ; A5D8U8 ; A6NFI4 ; D6W5X9
Molecular Weight	33,697 Da
NCBI Official Full Name	calponin-2 isoform a
NCBI Official Synonym Full Names	calponin 2
NCBI Official Symbol	CNN2
NCBI Protein Information	calponin-2; neutral calponin; calponin H2, smooth muscle
UniProt Protein Name	Calponin-2
UniProt Gene Name	CNN2

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/calponin-2-cnn2-antibody-western-blot-immunohistochemistry-flow-cytometry-elisa/126530>