

Rabbit anti-Human ATG4C Polyclonal Antibody | anti-ATG4C antibody

Anti-ATG4C Antibody Picoband

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA126561	Rabbit	Human	ELISA, Flow Cytometry, Functional Assay, Immunofluorescence, Immunocytochemistry, Western Blot

PRODUCT SYNONYMS

ATG4C; Polyclonal Antibody; Anti-ATG4C Antibody Picoband; anti-ATG4C antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Purity/Purification	Immunogen affinity purified.
Form/Format	Lyophilized Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml. (varies by lot)
Applicable Applications for anti-ATG4C antibody	ELISA, FCM/FACS (Flow Cytometry), IF (Immunofluorescence), ICC (Immunocytochemistry), WB (Western Blot)
Reconstitution	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Immunogen	E Coli-derived human ATG4C recombinant protein (Position: M1-L457).
Preparation and Storage	Store at -20 degree C for one year from date of receipt. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for six months.Avoid repeated freezing and thawing.

IMAGES



FCM/FACS (Flow Cytometry)

Figure 3. Flow Cytometry analysis of U251 cells using anti-ATG4C antibody (AAA126561).Overlay histogram showing U251 cells stained with AAA126561 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-ATG4C Antibody (AAA126561, 1 ug/1x10⁶ cells) for 30 min at 20 degree C. DyLight488 conjugated goat anti-rabbit IgG was used as secondary antibody for 30 minutes at 20 degree C. Isotype control antibody (Green line) was rabbit IgG (1 ug/1x10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.



IF (Immunofluorescence)

Figure 2. IF analysis of ATG4C using anti-ATG4C antibody (AAA126561) and anti-Tubulin Alpha antibody (M03989-3).ATG4C was detected in immunocytochemical section of U2OS cells. Enzyme antigen retrieval was performed using IHC enzyme antigen retrieval reagent (AR0022) for 15 mins. The cells were blocked with 10% goat serum. And then incubated with 5 ug/mL rabbit anti-ATG4C Antibody (AAA126561) and mouse anti-Tubulin Alpha antibody (M03989-3) overnight at 4 degree C. Cy3 Conjugated Goat Anti-Rabbit IgG (BA1032) and DyLight488 Conjugated Goat Anti-Mouse IgG were used as secondary antibody at 1:100 dilution and incubated for 30 minutes at 37 degree C. The section was counterstained with DAPI. Visualize using a fluorescence microscope and filter sets appropriate for the label used.



WB (Western Blot)

Figure 1. Western blot analysis of ATG4C using anti-ATG4C antibody (AAA126561). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel)/90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: human 293T whole cell lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-ATG4C antigen affinity purified polyclonal antibody (#AAA126561) at 0.5 ug/mL overnight at 4 degree C, then washed with TBS-0.1% Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for ATG4C at approximately 52 kDa. The expected band size for ATG4C is at 52 kDa.

ADDITIONAL INFORMATION

Related Product Information for anti-ATG4C antibody	Autophagy is the process by which endogenous proteins and damaged organelles are destroyed intracellularly. Autophagy is postulated to be essential for cell homeostasis and cell remodeling during differentiation, metamorphosis, non-apoptotic cell death, and aging. Reduced levels of autophagy have been described in some malignant tumors, and a role for autophagy in controlling the unregulated cell growth linked to cancer has been proposed. This gene encodes a member of the autophagin protein family. The encoded protein is also designated as a member of the C-54 family of cysteine proteases. Alternate transcriptional splice variants, encoding the same protein, have been characterized.
Product Categories/Family for anti-ATG4C antibody	Primary Antibodies, Rabbit Polyclonal Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	30410844
NCBI GeneID	84938
NCBI Accession #	NP_116241.2
NCBI GenBank Nucleotide #	NM_032852.3
UniProt Accession #	Q96DT6; Q96K04; A6NLR8; D3DQ58
Molecular Weight	52,497 Da
NCBI Official Full Name	cysteine protease ATG4C
NCBI Official Synonym Full Names	autophagy related 4C, cysteine peptidase
NCBI Official Symbol	ATG4C
NCBI Official Synonym Symbols	APG4C; AUTL1; AUTL3; APG4-C
NCBI Protein Information	cysteine protease ATG4C; autophagin-3; APG4 autophagy 4 homolog C; AUT-like 3 cysteine endopeptidase; ATG4 autophagy related 4 homolog C; AUT-like 1, cysteine endopeptidase; autophagy-related protein 4 homolog C; autophagy-related cysteine endopeptidase 3
UniProt Protein Name	Cysteine protease ATG4C
UniProt Gene Name	ATG4C
UniProt Synonym Gene Names	APG4C; AUTL1; AUTL3
UniProt Entry Name	ATG4C_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/atg4c-antibody-flow-cytometry-immunofluorescence-immunocytochemistry-western-blot-elisa/126561>