

Mouse anti-Human ALPP Monoclonal Antibody | anti-ALPP antibody

Anti-ALPP Antibody Picoband (monoclonal, 6C7G3)


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA126898	Mouse	Human	Flow Cytometry, Functional Assay, Immunofluorescence, Immunocytochemistry, Immunohistochemistry, Western Blot

PRODUCT SYNONYMS

ALPP; Monoclonal Antibody; Anti-ALPP Antibody Picoband (monoclonal; 6C7G3); anti-ALPP antibody


PRODUCT SPECIFICATIONS

Host	Mouse
Reactivity	Human
Clonality	Monoclonal
Isotype	Mouse IgG2b
Clone Number	6C7G3
Purity/Purification	Immunogen affinity purified.
Form/Format	Lyophilized Each vial contains 4 mg Trehalose, 0.9 mg NaCl and 0.2 mg Na ₂ HPO ₄ .
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml. (varies by lot)
Applicable Applications for anti-ALPP antibody	FCM/FACS (Flow Cytometry), IF (Immunofluorescence), ICC (Immunocytochemistry), IHC (Immunohistochemistry), WB (Western Blot)
Reconstitution	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Immunogen	A synthetic peptide corresponding to a sequence at the C-terminus of human ALPP (403-433aa KARDRKAYTVLLYGNGPGYVLKDGARPDVTE).
Preparation and Storage	Store at -20 degree C for one year from date of receipt. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for six months. Avoid repeated freezing and thawing.




FCM/FACS (Flow Cytometry)

Figure 5. Flow Cytometry analysis of SiHa cells using anti-ALPP antibody (AAA126898). Overlay histogram showing SiHa cells stained with AAA126898 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with mouse anti-ALPP Antibody (AAA126898, 1 ug/1x10⁶ cells) for 30 min at 20 degree C. DyLight488 conjugated goat anti-mouse IgG was used as secondary antibody for 30 minutes at 20 degree C. Isotype control antibody (Green line) was mouse IgG (1 ug/1x10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.




IHC (Immunohistochemisry)

Figure 3. IHC analysis of ALPP using anti-ALPP antibody (AAA126898).ALPP was detected in a paraffin-embedded section of human placenta tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml mouse anti-ALPP Antibody (AAA126898) overnight at 4 degree C. Peroxidase Conjugated Goat Anti-mouse IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using HRP Conjugated Mouse IgG Super Vision Assay Kit with DAB as the chromogen.




IF (Immunofluorescence)

Figure 4. IF analysis of ALPP using anti-ALPP antibody (AAA126898).ALPP was detected in an immunocytochemical section of HeLa cells. Enzyme antigen retrieval was performed using IHC enzyme antigen retrieval reagent (AR0022) for 15 mins. The cells were blocked with 10% goat serum. And then incubated with 5 ug/mL mouse anti-ALPP Antibody (AAA126898) overnight at 4 degree C. DyLight488 Conjugated Goat Anti-Mouse IgG was used as secondary antibody at 1:100 dilution and incubated for 30 minutes at 37 degree C. The section was counterstained with DAPI. Visualize using a fluorescence microscope and filter sets appropriate for the label used.



IF (Immunofluorescence)

Figure 2. IF analysis of ALPP using anti-ALPP antibody (AAA126898).ALPP was detected in a paraffin-embedded section of human placenta tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 5 ug/mL mouse anti-ALPP Antibody (AAA126898) overnight at 4 degree C. Biotin conjugated goat anti-mouse IgG (BA1001) was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using DyLight488 Conjugated Avidin (BA1128). The section was counterstained with DAPI. Visualize using a fluorescence microscope and filter sets appropriate for the label used.



WB (Western Blot)

Figure 1. Western blot analysis of ALPP using anti-ALPP antibody (AAA126898). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel)/90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: human placenta tissue lysates, Lane 2: human HepG2 whole cell lysates, Lane 3: human A431 whole cell lysates, Lane 4: human HeLa whole cell lysates, Lane 5: human SW620 whole cell lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with mouse anti-ALPP antigen affinity purified monoclonal antibody (#AAA126898) at 0.5 ug/mL overnight at 4 degree C, then washed with TBS-0.1% Tween 3 times with 5 minutes each and probed with a goat anti-mouse IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for ALPP at approximately 70 kDa. The expected band size for ALPP is at 70 kDa.

ADDITIONAL INFORMATION

Related Product Information for anti-ALPP antibody	Alkaline phosphatase, placental type also known as placental alkaline phosphatase (PLAP) is an allosteric enzyme that in humans is encoded by the ALPP gene. The protein encoded by this gene is an alkaline phosphatase, a metalloenzyme that catalyzes the hydrolysis of phosphoric acid monoesters. It belongs to a multigene family composed of four alkaline phosphatase isoenzymes. The enzyme functions as a homodimer and has a catalytic site containing one magnesium and two zinc ions, which are required for its enzymatic function. The protein is primarily expressed in placental and endometrial tissue; however, strong ectopic expression has been detected in ovarian adenocarcinoma, serous cystadenocarcinoma, and other ovarian cancer cells.
Product Categories/Family for anti-ALPP antibody	Primary Antibodies , Mouse Monoclonal Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	94721246
NCBI GeneID	250
NCBI Accession #	NP_001623.3
NCBI GenBank Nucleotide #	NM_001632.3
UniProt Accession #	P05187 ; P05188 ; P06861 ; Q53S78 ; Q96DB7
Molecular Weight	57,954 Da
NCBI Official Full Name	alkaline phosphatase, placental type preproprotein
NCBI Official Synonym Full Names	alkaline phosphatase, placental
NCBI Official Symbol	ALPP
NCBI Official Synonym Symbols	ALP; PALP; PLAP; PLAP-1
NCBI Protein Information	alkaline phosphatase, placental type; Regan isozyme; glycerophosphatase; alkaline phosphomonoesterase; placental alkaline phosphatase 1; alkaline phosphatase Regan isozyme

UniProt Protein Name	Alkaline phosphatase, placental type
UniProt Gene Name	ALPP
UniProt Synonym Gene Names	PLAP; PLAP-1
UniProt Entry Name	PPB1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/monoclonal/alpp-antibody-clone-6c7g3-flow-cytometry-immunofluorescence-immunocytochemistry-immunohistochemistry-western-blot-elisa/126898>