

Mouse anti-Human PC4/SUB1 Monoclonal Antibody | anti-SUB1 antibody

Anti-PC4/SUB1 Antibody Picoband (monoclonal, 2D13E3)


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA126915	Mouse	Human	Flow Cytometry, Functional Assay, Western Blot

PRODUCT SYNONYMS


PC4/SUB1; Monoclonal Antibody; Anti-PC4/SUB1 Antibody Picoband (monoclonal; 2D13E3); anti-SUB1 antibody

PRODUCT SPECIFICATIONS

Host	Mouse
Reactivity	Human
Clonality	Monoclonal
Isotype	Mouse IgG2b
Clone Number	2D13E3
Purity/Purification	Immunogen affinity purified.
Form/Format	Lyophilized Each vial contains 4 mg Trehalose, 0.9 mg NaCl and 0.2 mg Na ₂ HPO ₄ .
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml. (varies by lot)
Applicable Applications for anti-SUB1 antibody	FCM/FACS (Flow Cytometry), WB (Western Blot)
Reconstitution	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Immunogen	E Coli-derived human PC4/SUB1 recombinant protein (Position: N62-L127).
Preparation and Storage	Store at -20 degree C for one year from date of receipt. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for six months. Avoid repeated freezing and thawing.



FCM/FACS (Flow Cytometry)
Figure 2. Flow Cytometry analysis of PC-3 cells using anti-PC4/SUB1 antibody (AAA126915).Overlay histogram showing PC-3 cells stained with AAA126915 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with mouse anti-PC4/SUB1 Antibody (AAA126915, 1 ug/1x10⁶ cells) for 30 min at 20 degree C. DyLight488 conjugated goat anti-mouse IgG was used as secondary antibody for 30 minutes at 20 degree C. Isotype control antibody (Green line) was mouse IgG (1 ug/1x10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.



WB (Western Blot)
Figure 1. Western blot analysis of PC4/SUB1 using anti-PC4/SUB1 antibody (AAA126915).Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel)/90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.Lane 1: human HeLa whole cell lysates,Lane 2: human Jurkat whole cell lysates,Lane 3: human A549 whole cell lysates,Lane 4: human HepG2 whole cell lysates.After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with mouse anti-PC4/SUB1 antigen affinity purified monoclonal antibody (#AAA126915) at 0.5 ug/mL overnight at 4 degree C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-mouse IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for PC4/SUB1 at approximately 18 kDa. The expected band size for PC4/SUB1 is at 18 kDa.

ADDITIONAL INFORMATION

Related Product Information for anti-SUB1 antibody	Activated RNA polymerase II transcriptional coactivator p15, also known as positive cofactor 4 (PC4) or SUB1 homolog, is a protein that in humans is encoded by the SUB1 gene. This gene is mapped to 5p13.3. The transcriptional cofactor PC4 is an ancient single-strand DNA (ssDNA)-binding protein that has a homologue in bacteriophage T5 where it is likely the elusive replicative ssDNA-binding protein. The recombinant PC4 is shown to function identically to the native protein through its interaction with TAFs.
Product Categories/Family for anti-SUB1 antibody	Primary Antibodies, Mouse Monoclonal Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	1709514
NCBI GeneID	10923
UniProt Accession #	P53999; Q96L29
Molecular Weight	127
NCBI Official Full Name	Activated RNA polymerase II transcriptional coactivator p15
NCBI Official Synonym Full Names	SUB1 homolog (S. cerevisiae)
NCBI Official Symbol	SUB1
NCBI Official Synonym Symbols	P15; PC4; p14
NCBI Protein Information	activated RNA polymerase II transcriptional coactivator p15; positive cofactor 4; activated RNA polymerase II transcription cofactor 4
UniProt Protein Name	Activated RNA polymerase II transcriptional coactivator p15
UniProt Gene Name	SUB1

UniProt Synonym Gene Names	PC4; RPO2TC1; PC4
UniProt Entry Name	TCP4_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/monoclonal/pc4-sub1-antibody-clone-2d13e3-flow-cytometry-western-blot-elisa/126915>