

Mouse Integrin linked ILK Monoclonal Antibody | anti-ILK antibody

Anti-Integrin linked ILK Antibody Picoband (monoclonal, 3C7E1)


CATALOG # AAA126922	HOST Mouse	REACTIVITY Human, Mouse, Rat	APPLICATIONS Immunohistochemistry, Western Blot
-------------------------------	----------------------	--	---

PRODUCT SYNONYMS


Integrin linked ILK; Monoclonal Antibody; Anti-Integrin linked ILK Antibody Picoband (monoclonal; 3C7E1); anti-ILK antibody

PRODUCT SPECIFICATIONS


Host	Mouse
Reactivity	Human, Mouse, Rat
Clonality	Monoclonal
Isotype	Mouse IgG2b
Clone Number	3C7E1
Purity/Purification	Immunogen affinity purified.
Form/Format	Lyophilized Each vial contains 4 mg Trehalose, 0.9 mg NaCl and 0.2 mg Na ₂ HPO ₄ .
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml. (varies by lot)
Applicable Applications for anti-ILK antibody	IHC (Immunohistochemistry), WB (Western Blot)
Reconstitution	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Immunogen	E Coli-derived human Integrin linked ILK recombinant protein (Position: M1-K452).
Preparation and Storage	Store at -20 degree C for one year from date of receipt. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for six months. Avoid repeated freezing and thawing.




IHC (Immunohistochemistry)
Figure 4. IHC analysis of Integrin linked ILK using anti-Integrin linked ILK antibody (AAA126922).Integrin linked ILK was detected in a paraffin-embedded section of mouse kidney tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml mouse anti-Integrin linked ILK Antibody (AAA126922) overnight at 4 degree C. Peroxidase Conjugated Goat Anti-mouse IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using HRP Conjugated Mouse IgG Super Vision Assay Kit with DAB as the chromogen.



IHC (Immunohistochemistry)
Figure 3. IHC analysis of Integrin linked ILK using anti-Integrin linked ILK antibody (AAA126922).Integrin linked ILK was detected in a paraffin-embedded section of human spleen tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml mouse anti-Integrin linked ILK Antibody (AAA126922) overnight at 4 degree C. Peroxidase Conjugated Goat Anti-mouse IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using HRP Conjugated Mouse IgG Super Vision Assay Kit with DAB as the chromogen.



IHC (Immunohistochemistry)
Figure 2. IHC analysis of Integrin linked ILK using anti-Integrin linked ILK antibody (AAA126922).Integrin linked ILK was detected in a paraffin-embedded section of human squamous cell lung carcinoma tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml mouse anti-Integrin linked ILK Antibody (AAA126922) overnight at 4 degree C. Peroxidase Conjugated Goat Anti-mouse IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using HRP Conjugated Mouse IgG Super Vision Assay Kit with DAB as the chromogen.



WB (Western Blot)
Figure 1. Western blot analysis of Integrin linked ILK using anti-Integrin linked ILK antibody (AAA126922).Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel)/90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.Lane 1: human MCF-7 whole cell lysates,Lane 2: human HeLa whole cell lysates,Lane 3: human U87 whole cell lysates,Lane 4: rat heart tissue lysates.Lane 5: rat kidney tissue lysates.Lane 6: mouse heart tissue lysates.Lane 7: mouse kidney tissue lysates.After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with mouse anti-Integrin linked ILK antigen affinity purified monoclonal antibody (#AAA126922) at 0.5 ug/mL overnight at 4 degree C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-mouse IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for Integrin linked ILK at approximately 51 kDa. The expected band size for Integrin linked ILK is at 51 kDa.

ADDITIONAL INFORMATION

Related Product Information for anti-ILK antibody

ILK, also known as Integrin-linked kinase, is a serine-threonine protein kinase. Transduction of extracellular matrix signals through integrins influences intracellular and extracellular functions, and appears to require interaction of integrin cytoplasmic domains with cellular proteins. Integrin-linked kinase (ILK) interacts with the cytoplasmic domain of beta-1 integrin. This gene was initially described to encode a serine/ threonine protein kinase with 4 ankyrin-like repeats, which associates with the cytoplasmic domain of beta integrins and acts as a proximal receptor kinase regulating integrin-mediated signal transduction. Multiple alternatively spliced transcript variants encoding the same protein have been found for this gene. Recent results showed that ILK contains 5 ankyrin-like repeats, and that the C-terminal kinase domain is actually a pseudo-kinase with adaptor function.

Product Categories/Family for anti-ILK antibody

Primary Antibodies, Mouse Monoclonal Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	62420873
NCBI GeneID	3611
NCBI Accession #	NP_001014794.1
NCBI GenBank Nucleotide #	NM_001014794.2
UniProt Accession #	Q13418; P57043; Q68DZ3; D3DQU0
Molecular Weight	51,419 Da
NCBI Official Full Name	integrin-linked protein kinase isoform 1
NCBI Official Synonym Full Names	integrin-linked kinase
NCBI Official Symbol	ILK
NCBI Official Synonym Symbols	P59; ILK-1; ILK-2; p59ILK
NCBI Protein Information	integrin-linked protein kinase; integrin-linked kinase-2; 59 kDa serine/threonine-protein kinase
UniProt Protein Name	Integrin-linked protein kinase
UniProt Gene Name	ILK
UniProt Synonym Gene Names	ILK1; ILK2
UniProt Entry Name	ILK_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/monoclonal/integrin-linked-ilk-antibody-clone-3c7e1-immunohistochemistry-western-blot-elisa/126922>