

Mouse anti-Human MASPIN Monoclonal Antibody | anti-SERPINB5 antibody

Anti-MASPIN Antibody Picoband (monoclonal, 7G4E1)


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA126926	Mouse	Human	Flow Cytometry, Functional Assay, Immunohistochemistry, Western Blot

PRODUCT SYNONYMS

MASPIN; Monoclonal Antibody; Anti-MASPIN Antibody Picoband (monoclonal; 7G4E1); anti-SERPINB5 antibody


PRODUCT SPECIFICATIONS

Host	Mouse
Reactivity	Human
Clonality	Monoclonal
Isotype	Mouse IgG2a
Clone Number	7G4E1
Purity/Purification	Immunogen affinity purified.
Form/Format	Lyophilized Each vial contains 4 mg Trehalose, 0.9 mg NaCl and 0.2 mg Na ₂ HPO ₄ .
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml. (varies by lot)
Applicable Applications for anti-SERPINB5 antibody	FCM/FACS (Flow Cytometry), IHC (Immunohistochemistry), WB (Western Blot)
Reconstitution	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Immunogen	E Coli-derived human MASPIN recombinant protein (Position: M1-A350). Human MASPIN shares 88% and 89% amino acid (aa) sequence identity with mouse and rat MASPIN, respectively.
Preparation and Storage	Store at -20 degree C for one year from date of receipt. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for six months. Avoid repeated freezing and thawing.




FCM/FACS (Flow Cytometry)

Figure 4. Flow Cytometry analysis of SiHa cells using anti-MASPIN antibody (AAA126926).Overlay histogram showing SiHa cells stained with AAA126926 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with mouse anti-MASPIN Antibody (AAA126926, 1 ug/1x10⁶ cells) for 30 min at 20 degree C. DyLight488 conjugated goat anti-mouse IgG was used as secondary antibody for 30 minutes at 20 degree C. Isotype control antibody (Green line) was mouse IgG (1 ug/1x10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.




IHC (Immunohistochemistry)

Figure 2. IHC analysis of MASPIN using anti-MASPIN antibody (AAA126926).MASPIN was detected in a paraffin-embedded section of human tonsil tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml mouse anti-MASPIN Antibody (AAA126926) overnight at 4 degree C. Peroxidase Conjugated Goat Anti-mouse IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using HRP Conjugated Mouse IgG Super Vision Assay Kit with DAB as the chromogen.



IHC (Immunohistochemistry)

Figure 3. IHC analysis of MASPIN using anti-MASPIN antibody (AAA126926).MASPIN was detected in a paraffin-embedded section of human laryngeal squamous cell carcinoma tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml mouse anti-MASPIN Antibody (AAA126926) overnight at 4 degree C. Peroxidase Conjugated Goat Anti-mouse IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using HRP Conjugated Mouse IgG Super Vision Assay Kit with DAB as the chromogen.



WB (Western Blot)

Figure 1. Western blot analysis of MASPIN using anti-MASPIN antibody (AAA126926).Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel)/90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.Lane 1: human HeLa whole cell lysates,Lane 2: human Hacat whole cell lysates,Lane 3: human Caco-2 whole cell lysates.After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with mouse anti-MASPIN antigen affinity purified monoclonal antibody (#AAA126926) at 0.5 ug/mL overnight at 4 degree C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-mouse IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for MASPIN at approximately 42 kDa. The expected band size for MASPIN is at 42 kDa.

ADDITIONAL INFORMATION

Related Product Information for anti-SERPINB5 antibody	SERPINB5 is also known as PI5 or maspin. Maspin (mammary serine protease inhibitor) is a protein that in humans is encoded by the SERPINB5 gene. Maspin is expressed in the skin, prostate, testis, intestine, tongue, lung, and the thymus. Maspin is a member of the serpin superfamily of serine protease inhibitors.[1] The primary function of most members of this family is to regulate the breakdown of proteins by inhibiting the catalytic activity of proteinases. Through this mechanism of action, serpins regulate a number of cellular processes includingphagocytosis, coagulation, and fibrinolysis.
Product Categories/Family for anti-SERPINB5 antibody	Primary Antibodies, Mouse Monoclonal Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	167860126
NCBI GeneID	5268
NCBI Accession #	NP_002630.2
NCBI GenBank Nucleotide #	NM_002639.4
UniProt Accession #	P36952; Q6N0B4; Q8WW89; B2R6Y4
Molecular Weight	375
NCBI Official Full Name	serpin B5
NCBI Official Synonym Full Names	serpin peptidase inhibitor, clade B (ovalbumin), member 5
NCBI Official Symbol	SERPINB5
NCBI Official Synonym Symbols	PI5; maspin
NCBI Protein Information	serpin B5; PI-5; peptidase inhibitor 5; protease inhibitor 5 (maspin); serine (or cysteine) proteinase inhibitor, clade B (ovalbumin), member 5
UniProt Protein Name	Serpin B5
UniProt Gene Name	SERPINB5
UniProt Synonym Gene Names	PI5; PI-5
UniProt Entry Name	SPB5_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/monoclonal/maspin-antibody-clone-7g4e1-flow-cytometry-immunohistochemistry-western-blot-elisa/126926>