

Rabbit PKC Alpha/PRKCA Polyclonal Antibody | anti-PKCApha/PRKCA antibody

Anti-PKC Alpha/PRKCA Antibody Picoband


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA127009	Rabbit	Human, Mouse, Rat	ELISA, Flow Cytometry, Functional Assay, Western Blot

PRODUCT SYNONYMS

PKC Alpha/PRKCA; Polyclonal Antibody; Anti-PKC Alpha/PRKCA Antibody Picoband; Argininosuccinate lyase; ASAL; 4.3.2.1; Arginosuccinase; ASL; anti-PKCApha/PRKCA antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	Rabbit IgG
Purity/Purification	Immunogen affinity purified.
Form/Format	Lyophilized. Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Applicable Applications for anti-PKCApha/PRKCA antibody	ELISA, FCM/FACS (Flow Cytometry), WB (Western Blot)
Immunogen	E coli-derived human PKC Alpha/PRKCA recombinant protein (Position: Q142-V672).
Subcellular Localization	Extracellular matrix
Reconstitution	Adding 0.2 ml of distilled water will yield a concentration of 500ug/ml.
Cross Reactivity	No cross-reactivity with other proteins.
Protein Function	Cleaves collagens of types I, II, and III at one site in the helical domain. Also cleaves collagens of types VII and X. In case of HIV infection, interacts and cleaves the secreted viral Tat protein, leading to a decrease in neuronal Tat's mediated neurotoxicity.
Preparation and Storage	At -20 degree C for one year from date of receipt. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for six months.Avoid repeated freezing and thawing.



FCM/FACS (Flow Cytometry)

Figure 2. Flow Cytometry analysis of SiHa cells using anti-PKC Alpha/PRKCA antibody (AAA127009).Overlay histogram showing SiHa cells stained with AAA127009 (Blue line). To facilitate intracellular staining, cells were fixed with 4% paraformaldehyde and permeabilized with permeabilization buffer. The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-PKC Alpha/PRKCA Antibody (AAA127009, 1ug/1x106 cells) for 30 min at 20 degree C. DyLight488 conjugated goat anti-rabbit IgG was used as secondary antibody for 30 minutes at 20 degree C. Isotype control antibody (Green line) was rabbit IgG (1ug/1x106) used under the same conditions. Unlabelled sample (Red line) was also used as a control.



WB (Western Blot)

Figure 1. Western blot analysis of PKC Alpha/PRKCA using anti-PKC Alpha/PRKCA antibody (AAA127009).Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel)/90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.Lane 1: human Jurkat whole cell lysates,Lane 2: human 293T cell lysates,Lane 3: human HeLa whole cell lysates,Lane 4: human K562 whole cell lysates,Lane 5: rat C6 whole cell lysates,Lane 6: mouse NIH/3T3 whole cell lysates.After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-PKC Alpha/PRKCA antigen affinity purified polyclonal antibody (#AAA127009) at 0.5ug/mL overnight at 4 degree C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for PKC Alpha/PRKCA at approximately 80 kDa. The expected band size for PKC Alpha/PRKCA is at 77 kDa.

ADDITIONAL INFORMATION

Related Product Information for anti-PKCAAlpha/PRKCA antibody	The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Product Categories/Family for anti-PKCAAlpha/PRKCA antibody	Primary Antibodies; Cancer; Energy Metabolism; Energy Transfer Pathways; Metabolic Signaling Pathways; Metabolism; Pathways and Processes; Signal Transduction
References	1. Braz, J. C.; Gregory, K.; Pathak, A.; Zhao, W.; Sahin, B.; Klevitsky, R.; Kimball, T. F.; Lorenz, J. N.; Nairn, A. C.; Liggett, S. B.; Bodi, I.; Wang, S.; and 9 others : PKC-alpha regulates cardiac contractility and propensity toward heart failure. Nature Med. 10: 248-254, 2004.2. Latos-Bielenska, A.; Klett, C.; Just, W.; Hameister, H. : Refinement of localization of the human genes for myeloperoxidase (MPO), protein kinase C, alpha polypeptide, PRKCA, and the DNA fragment D17S21 on chromosome 17q. Hereditas 115: 69-72, 1991.

NCBI AND UNIPROT INFORMATION

NCBI GI #	4506067
NCBI GeneID	5578
NCBI Accession #	NP_002728.1
NCBI GenBank Nucleotide #	NM_002737.2
UniProt Accession #	P17252; Q15137; Q32M72; Q96RE4; B5BU22
Molecular Weight	76,750 Da
NCBI Official Full Name	protein kinase C alpha type
NCBI Official Synonym Full Names	protein kinase C, alpha
NCBI Official Symbol	PRKCA
NCBI Official Synonym Symbols	AAG6; PKCA; PRKACA; PKC-alpha

NCBI Protein Information	protein kinase C alpha type; PKC-A; aging-associated gene 6
UniProt Protein Name	Protein kinase C alpha type
UniProt Gene Name	PRKCA
UniProt Synonym Gene Names	PKCA; PRKACA; PKC-A; PKC-alpha
UniProt Entry Name	KPCA_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/pkc-alpha-prkca-antibody-flow-cytometry-western-blot-elisa/127009>