

Rabbit anti-Human ABCC3 Polyclonal Antibody | anti-ABCC3 antibody

Anti-ABCC3 Antibody Picoband


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA127116	Rabbit	Human	Flow Cytometry, Functional Assay, Western Blot, ELISA

PRODUCT SYNONYMS


ABCC3; Polyclonal Antibody; Anti-ABCC3 Antibody Picoband; BAG family molecular chaperone regulator 1; Fc fragment of IgG receptor Ia; High affinity immunoglobulin gamma Fc receptor I; IgG Fc receptor I; Fc-gamma RI; FcRI; Fc-gamma RIA; FcgammaRIa; CD64; FCGR1A; FCG1; FCGR1; IGFR1; anti-ABCC3 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Monocyte/macrophage specific.
Purity/Purification	Immunogen affinity purified.
Form/Format	Lyophilized. Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na2HPO4.
Applicable Applications for anti-ABCC3 antibody	FCM/FACS (Flow Cytometry), WB (Western Blot), ELISA
Immunogen	E coli-derived human ABCC3 recombinant protein (Position: H22-G1351).
Subcellular Localization	Cell membrane
Reconstitution	Adding 0.2 ml of distilled water will yield a concentration of 500ug/ml.
Cross Reactivity	No cross-reactivity with other proteins.
Protein Function	High affinity receptor for the Fc region of immunoglobulins gamma. Functions in both innate and adaptive immune responses.
Preparation and Storage	At -20 degree C for one year from date of receipt. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for six months.Avoid repeated freezing and thawing.



FCM/FACS (Flow Cytometry)
Figure 2. Flow Cytometry analysis of HepG2 cells using anti-ABCC3 antibody (AAA127116). Overlay histogram showing HepG2 cells stained with AAA127116 (Blue line). The cells were fixed with 4% paraformaldehyde and blocked with 10% normal goat serum. And then incubated with rabbit anti-ABCC3 Antibody (AAA127116, 1ug/1x10⁶ cells) for 30 min at 20 degree C. DyLight488 conjugated goat anti-rabbit IgG was used as secondary antibody for 30 minutes at 20 degree C. Isotype control antibody (Green line) was rabbit IgG (1ug/1x10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.



WB (Western Blot)
Figure 1. Western blot analysis of ABCC3 using anti-ABCC3 antibody (AAA127116). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel)/90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: human HepG2 whole cell lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-ABCC3 antigen affinity purified polyclonal antibody (#AAA127116) at 0.5ug/mL overnight at 4 degree C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for ABCC3 at approximately 200 kDa. The expected band size for ABCC3 is at 200 kDa.

ADDITIONAL INFORMATION

Related Product Information for anti-ABCC3 antibody	The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Product Categories/Family for anti-ABCC3 antibody	Primary Antibodies; Immunoglobulins; Immunology; Innate Immunity
References	1. Belinsky, M. G., Bain, L. J., Balsara, B. B., Testa, J. R., Kruh, G. D. Characterization of MOAT-C and MOAT-D, new members of the MRP/cMOAT subfamily of transporter proteins. J. Nat. Cancer Inst. 90: 1735-1741, 1998. 2. Fromm, M. F., Leake, B., Roden, D. M., Wilkinson, G. R., Kim, R. B. Human MRP3 transporter: identification of the 5-prime flanking region, genomic organization and alternative splice variants. Biochim. Biophys. Acta 1415: 369-374, 1999. 3. Kiuchi, Y., Suzuki, H., Hirohashi, T., Tyson, C. A., Sugiyama, Y. cDNA cloning and inducible expression of human multidrug resistance associated protein 3 (MRP3). FEBS Lett. 433: 149-152,1998.

NCBI AND UNIPROT INFORMATION

NCBI GI #	221316556
NCBI GeneID	8714
NCBI Accession #	NP_001137542.1
NCBI GenBank Nucleotide #	NM_001144070.1
UniProt Accession #	O15438; O60265; O60922; O75621; O95078; O95289; O95290; Q86X85; Q9UN52; B2RPA9; D3DTX9
Molecular Weight	169,343 Da
NCBI Official Full Name	canalicular multispecific organic anion transporter 2 isoform 2
NCBI Official Synonym Full Names	ATP-binding cassette, sub-family C (CFTR/MRP), member 3
NCBI Official Symbol	ABCC3
NCBI Official Synonym Symbols	MLP2; MRP3; ABC31; MOAT-D; cMOAT2; EST90757

NCBI Protein Information	canalicular multispecific organic anion transporter 2; multidrug resistance associated protein; multidrug resistance-associated protein 3; ATP-binding cassette sub-family C member 3; multi-specific organic anion transporter D; canicular multispecific orga
UniProt Protein Name	Canalicular multispecific organic anion transporter 2
UniProt Gene Name	ABCC3
UniProt Synonym Gene Names	CMOAT2; MLP2; MRP3; MOAT-D
UniProt Entry Name	MRP3_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/abcc3-antibody-flow-cytometry-western-blot-elisa/127116>