

Rabbit anti-Human PEG3 Polyclonal Antibody | anti-PEG3 antibody

Anti-PEG3 Antibody Picoband


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA127125	Rabbit	Human	ELISA, Flow Cytometry, Functional Assay, Immunofluorescence, Immunocytochemistry, Western Blot

PRODUCT SYNONYMS

PEG3; Polyclonal Antibody; Anti-PEG3 Antibody Picoband; Serine/threonine-protein kinase AtPK1/AtPK6; 70 kDa ribosomal protein S6 kinase 1 antibody; KS6B1_HUMAN antibody; p70 alpha antibody; P70 beta 1 antibody; p70 ribosomal S6 kinase alpha antibody; p70 ribosomal S6 kinase beta 1 antibody; p70 S6 kinase alpha antibody; P70 S6 Kinase antibody; p70 S6 kinase alpha 1 antibody; p70 S6 kinase alpha 2 antibody; p70 S6K antibody; p70 S6K-alpha antibody; p70 S6KA antibody; p70(S6K) alpha antibody; p70(S6K)-alpha antibody; p70-alpha antibody; p70-S6K 1 antibody; p70-S6K antibody; P70S6K antibody; P70S6K1 antibody; p70S6Kb antibody; PS6K antibody; Ribosomal protein S6 kinase 70kDa polypeptide 1 antibody; Ribosomal protein S6 kinase beta 1 antibody; Ribosomal protein S6 kinase beta-1 antibody; Ribosomal protein S6 kinase I antibody; RPS6KB1 antibody; S6K antibody; S6K-beta-1 antibody; S6K1 antibody; Serine/threonine kinase 14 alpha antibody; Serine/threonine-protein kinase 14A antibody; STK14A antibody; anti-PEG3 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Specificity	Expressed in all tissues.
Purity/Purification	Immunogen affinity purified.
Form/Format	Lyophilized. Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na2HPO4.
Applicable Applications for anti-PEG3 antibody	ELISA, FCM/FACS (Flow Cytometry), IF (Immunofluorescence), ICC (Immunocytochemistry), WB (Western Blot)
Immunogen	E coli-derived human PEG3 recombinant protein (Position: M1-R1548). Human PEG3 shares 62.3% amino acid (aa) sequence identity with mouse PEG3.
Subcellular Localization	Cytoplasm, Nucleus
Reconstitution	Adding 0.2 ml of distilled water will yield a concentration of 500ug/ml.
Cross Reactivity	No cross reactivity with other proteins.
Protein Function	Downstream effector of TOR signaling pathway involved in osmotic stress response. Could be involved in the control of plant growth and development. Phosphorylates the ribosomal proteins P14, P16 and S6. Functions as a repressor of cell proliferation and required for maintenance of chromosome stability and ploidy levels through the RBR1-E2F pathway.
Preparation and Storage	At -20 degree C for one year from date of receipt. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for six months.Avoid repeated freezing and thawing.




FCM/FACS (Flow Cytometry)

Figure 3. Flow Cytometry analysis of PC-3 cells using anti-PEG3 antibody (AAA127125). Overlay histogram showing PC-3 cells stained with AAA127125 (Blue line). To facilitate intracellular staining, cells were fixed with 4% paraformaldehyde and permeabilized with permeabilization buffer. The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-PEG3 Antibody (AAA127125, 1ug/1x10⁶ cells) for 30 min at 20 degree C. DyLight488 conjugated goat anti-rabbit IgG was used as secondary antibody for 30 minutes at 20 degree C. Isotype control antibody (Green line) was rabbit IgG (1ug/1x10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.



IF (Immunofluorescence)

Figure 2. IF analysis of PEG3 using anti-PEG3 antibody (AAA127125). PEG3 was detected in an immunocytochemical section of U2OS cells. Enzyme antigen retrieval was performed using IHC enzyme antigen retrieval reagent for 15 mins. The cells were blocked with 10% goat serum. And then incubated with 5ug/mL rabbit anti-PEG3 Antibody (AAA127125) overnight at 4 degree C. Cy3 Conjugated Goat Anti-Rabbit IgG (BA1032) was used as secondary antibody at 1:100 dilution and incubated for 30 minutes at 37 degree C. The tissue section was developed using Phalloidin-iFluor 488 Conjugated. Visualize using a fluorescence microscope and filter sets appropriate for the label used.



WB (Western Blot)

Figure 1. Western blot analysis of PEG3 using anti-PEG3 antibody (AAA127125). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel)/90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: human HepG2 whole cell lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-PEG3 antigen affinity purified polyclonal antibody (#AAA127125) at 0.5ug/mL overnight at 4 degree C, then washed with TBS-0.1% Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for PEG3 at approximately 200 kDa. The expected band size for PEG3 is at 181 kDa.

ADDITIONAL INFORMATION

Related Product Information for anti-PEG3 antibody	The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Product Categories/Family for anti-PEG3 antibody	Primary Antibodies; Signal Transduction; Signaling Pathway; Small G Proteins

References	1. Biankin, A. V., Waddell, N., Kassahn, K. S., Gingras, M.-C., Muthuswamy, L. B., Johns, A. L., Miller, D. K., Wilson, P. J., Patch, A.-M., Wu, J., Chang, D. K., Cowley, M. J., and 116 others. Pancreatic cancer genomes reveal aberrations in axon guidance pathway genes. <i>Nature</i> 491: 399-405, 2012. 2. Dorus, S., Vallender, E. J., Evans, P. D., Anderson, J. R., Gilbert, S. L., Mahowald, M., Wyckoff, G. J., Malcom, C. M., Lahn, B. T. Accelerated evolution of nervous system genes in the origin of Homo sapiens. <i>Cell</i> 119: 1027-1040, 2004. 3. Hiby, S. E., Lough, M., Keverne, E. B., Surani, M. A., Loke, Y. W., King, A. Paternal monoallelic expression of PEG3 in the human placenta. <i>Hum. Molec. Genet.</i> 10: 1093-1100, 2001.
------------	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	226053126
NCBI GeneID	5178
NCBI Accession #	NP_001139656.1
NCBI GenBank Nucleotide #	NM_001146184.1
UniProt Accession #	Q9GZU2; P78418; Q5H9P9; Q7Z7H7; Q8TF75; Q9GZY2; A7E2B8; B4DIM4; C9JP50
Molecular Weight	180,827 Da
NCBI Official Full Name	paternally-expressed gene 3 protein isoform 1
NCBI Official Synonym Full Names	paternally expressed 3
NCBI Official Symbol	PEG3
NCBI Official Synonym Symbols	PW1; ZNF904; ZSCAN24; ZKSCAN22
NCBI Protein Information	paternally-expressed gene 3 protein; Kruppel-type zinc finger protein; zinc finger and SCAN domain-containing protein 24
UniProt Protein Name	Paternally-expressed gene 3 protein
UniProt Gene Name	PEG3
UniProt Synonym Gene Names	KIAA0287; ZSCAN24
UniProt Entry Name	PEG3_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.