

Rabbit PMM2 Polyclonal Antibody | anti-PMM2 antibody

Anti-PMM2 Antibody Picoband


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA127163	Rabbit	Human, Mouse, Rat	ELISA, Flow Cytometry, Functional Assay, Immunofluorescence, Immunocytochemistry, Western Blot

PRODUCT SYNONYMS

PMM2; Polyclonal Antibody; Anti-PMM2 Antibody Picoband; Serine/threonine-protein kinase AtPK1/AtPK6; 70 kDa ribosomal protein S6 kinase 1 antibody; KS6B1_HUMAN antibody; p70 alpha antibody; P70 beta 1 antibody; p70 ribosomal S6 kinase alpha antibody; p70 ribosomal S6 kinase beta 1 antibody; p70 S6 kinase alpha antibody; P70 S6 Kinase antibody; p70 S6 kinase alpha 1 antibody; p70 S6 kinase alpha 2 antibody; p70 S6K antibody; p70 S6K-alpha antibody; p70 S6KA antibody; p70(S6K) alpha antibody; p70(S6K)-alpha antibody; p70-alpha antibody; p70-S6K 1 antibody; p70-S6K antibody; P70S6K antibody; P70S6K1 antibody; p70S6Kb antibody; PS6K antibody; Ribosomal protein S6 kinase 70kDa polypeptide 1 antibody; Ribosomal protein S6 kinase beta 1 antibody; Ribosomal protein S6 kinase beta-1 antibody; Ribosomal protein S6 kinase I antibody; RPS6KB1 antibody; S6K antibody; S6K-beta-1 antibody; S6K1 antibody; Serine/threonine kinase 14 alpha antibody; Serine/threonine-protein kinase 14A antibody; STK14A antibody; anti-PMM2 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Specificity	Expressed in all tissues.
Purity/Purification	Immunogen affinity purified.
Form/Format	Lyophilized. Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na2HPO4.
Applicable Applications for anti-PMM2 antibody	ELISA, FCM/FACS (Flow Cytometry), IF (Immunofluorescence), ICC (Immunocytochemistry), WB (Western Blot)
Immunogen	E coli-derived human PMM2 recombinant protein (Position: D48-S246). Human PMM2 shares 92.4% amino acid (aa) sequence identity with mouse PMM2.
Subcellular Localization	Cytoplasm, Nucleus
Reconstitution	Adding 0.2 ml of distilled water will yield a concentration of 500ug/ml.
Cross Reactivity	No cross reactivity with other proteins.
Protein Function	Downstream effector of TOR signaling pathway involved in osmotic stress response. Could be involved in the control of plant growth and development. Phosphorylates the ribosomal proteins P14, P16 and S6. Functions as a repressor of cell proliferation and required for maintenance of chromosome stability and ploidy levels through the RBR1-E2F pathway.
Preparation and Storage	At -20 degree C for one year from date of receipt. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for six months.Avoid repeated freezing and thawing.




FCM/FACS (Flow Cytometry)

Figure 3. Flow Cytometry analysis of HEL cells using anti-PMM2 antibody (AAA127163). Overlay histogram showing HEL cells stained with AAA127163 (Blue line). To facilitate intracellular staining, cells were fixed with 4% paraformaldehyde and permeabilized with permeabilization buffer. The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-PMM2 Antibody (AAA127163, 1ug/1x10⁶ cells) for 30 min at 20 degree C. DyLight488 conjugated goat anti-rabbit IgG was used as secondary antibody for 30 minutes at 20 degree C. Isotype control antibody (Green line) was rabbit IgG (1ug/1x10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.



IF (Immunofluorescence)

Figure 2. IF analysis of PMM2 using anti-PMM2 antibody (AAA127163) and anti-Beta Tubulin antibody (M01857-3). PMM2 was detected in immunocytochemical section of HELA cell. Enzyme antigen retrieval was performed using IHC enzyme antigen retrieval reagent for 15 mins. The cells were blocked with 10% goat serum. And then incubated with 5ug/mL rabbit anti-PMM2 Antibody (AAA127163) and mouse anti-Beta Tubulin antibody (M01857-3) overnight at 4 degree C. DyLight488 Conjugated Goat Anti-Rabbit IgG and Cy3 Conjugated Goat Anti-Mouse IgG (BA1031) were used as secondary antibody at 1:500 dilution and incubated for 30 minutes at 37 degree C. Visualize using a fluorescence microscope and filter sets appropriate for the label used.



WB (Western Blot)

Figure 1. Western blot analysis of PMM2 using anti-PMM2 antibody (AAA127163). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel)/90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: human HepG2 whole cell lysates, Lane 2: human A549 whole cell lysates, Lane 3: human MCF-7 whole cell lysates, Lane 4: rat small intestines tissue lysates, Lane 5: mouse small intestines tissue lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-PMM2 antigen affinity purified polyclonal antibody (#AAA127163) at 0.5ug/mL overnight at 4 degree C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for PMM2 at approximately 28 kDa. The expected band size for PMM2 is at 28 kDa.

ADDITIONAL INFORMATION

Related Product Information for anti-PMM2 antibody	The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Product Categories/Family for anti-PMM2 antibody	Primary Antibodies; Signal Transduction; Signaling Pathway; Small G Proteins

References	1. Bjursell, C., Erlandson, A., Nordling, M., Nilsson, S., Wahlstrom, J., Stibler, H., Kristiansson, B., Martinsson, T. PMM2 mutation spectrum, including 10 novel mutations, in a large CDG type 1A family material with a focus on Scandinavian families. Hum. Mutat. 16: 395-400, 2000.2. Bjursell, C., Wahlstrom, J., Berg, K., Stibler, H., Kristiansson, B., Matthijs, G., Martinsson, T. Detailed mapping of the phosphomannomutase 2 (PMM2) gene and mutation detection enable improved analysis for Scandinavian CDG type I families. Europ. J. Hum. Genet. 6: 603-611, 1998. 3. Bohles, H., Sewell, A. C., Gebhardt, B., Reinecke-Luthge, A., Kloppel, G., Marquardt, T. Hyperinsulinaemic hypoglycaemia: leading symptom in a patient with congenital disorder of glycosylation Ia (phosphomannomutase deficiency). J. Inherit. Metab. Dis. 24: 858-862, 2001.
------------	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	3024413
NCBI GeneID	5373
UniProt Accession #	O15305; A8K672; B7Z6R0; D3DUF3
Molecular Weight	246
NCBI Official Full Name	Phosphomannomutase 2
NCBI Official Synonym Full Names	phosphomannomutase 2
NCBI Official Symbol	PMM2
NCBI Official Synonym Symbols	PMI; CDG1; CDGS; PMI1; CDG1a; PMM 2
NCBI Protein Information	phosphomannomutase 2; phosphomannose isomerase 1
UniProt Protein Name	Phosphomannomutase 2
UniProt Gene Name	PMM2
UniProt Synonym Gene Names	PMM 2
UniProt Entry Name	PMM2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
 Online datasheet: <https://stage.aaabiotech.com/polyclonal/pmm2-antibody-flow-cytometry-immunofluorescence-immunocytochemistry-western-blot-elisa/127163>