

Rabbit anti-Mouse, Rat Osmr Polyclonal Antibody | anti-Osmr antibody

Anti-Osmr Antibody Picoband



CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA127212	Rabbit	Mouse, Rat	Flow Cytometry, Functional Assay, Western Blot, ELISA

PRODUCT SYNONYMS

Osmr; Polyclonal Antibody; Anti-Osmr Antibody Picoband; Serine/threonine-protein kinase AtPK1/AtPK6; 70 kDa ribosomal protein S6 kinase 1 antibody; KS6B1_HUMAN antibody; p70 alpha antibody; P70 beta 1 antibody; p70 ribosomal S6 kinase alpha antibody; p70 ribosomal S6 kinase beta 1 antibody; p70 S6 kinase alpha antibody; P70 S6 Kinase antibody; p70 S6 kinase alpha 1 antibody; p70 S6 kinase alpha 2 antibody; p70 S6K antibody; p70 S6K-alpha antibody; p70 S6KA antibody; p70(S6K) alpha antibody; p70(S6K)-alpha antibody; p70-alpha antibody; p70-S6K 1 antibody; p70-S6K antibody; P70S6K antibody; P70S6K1 antibody; p70S6Kb antibody; PS6K antibody; Ribosomal protein S6 kinase 70kDa polypeptide 1 antibody; Ribosomal protein S6 kinase beta 1 antibody; Ribosomal protein S6 kinase beta-1 antibody; Ribosomal protein S6 kinase I antibody; RPS6KB1 antibody; S6K antibody; S6K-beta-1 antibody; S6K1 antibody; Serine/threonine kinase 14 alpha antibody; Serine/threonine-protein kinase 14A antibody; STK14A antibody; anti-Osmr antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Specificity	Expressed in all tissues.
Purity/Purification	Immunogen affinity purified.
Form/Format	Lyophilized. Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na2HPO4.
Applicable Applications for anti-Osmr antibody	FCM/FACS (Flow Cytometry), WB (Western Blot), ELISA
Immunogen	E coli-derived mouse Osmr recombinant protein (Position: E673-Q968). Mouse Osmr shares 57.7% and 86.9% amino acid (aa) sequence identity with human and rat Osmr, respectively.
Subcellular Localization	Cytoplasm, Nucleus
Reconstitution	Adding 0.2 ml of distilled water will yield a concentration of 500ug/ml.
Cross Reactivity	No cross reactivity with other proteins.
Protein Function	Downstream effector of TOR signaling pathway involved in osmotic stress response. Could be involved in the control of plant growth and development. Phosphorylates the ribosomal proteins P14, P16 and S6. Functions as a repressor of cell proliferation and required for maintenance of chromosome stability and ploidy levels through the RBR1-E2F pathway.
Preparation and Storage	At -20 degree C for one year from date of receipt. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for six months.Avoid repeated freezing and thawing.



FCM/FACS (Flow Cytometry)
Figure 2. Flow Cytometry analysis of C2C12 cells using anti-Osmr antibody (AAA127212). Overlay histogram showing C2C12 cells stained with AAA127212 (Blue line). The cells were fixed with 4% paraformaldehyde and blocked with 10% normal goat serum. And then incubated with rabbit anti-Osmr Antibody (AAA127212, 1ug/1x106 cells) for 30 min at 20 degree C. DyLight488 conjugated goat anti-rabbit IgG was used as secondary antibody for 30 minutes at 20 degree C. Isotype control antibody (Green line) was rabbit IgG (1ug/1x106) used under the same conditions. Unlabelled sample (Red line) was also used as a control.

WB (Western Blot)
Figure 1. Western blot analysis of Osmr using anti-Osmr antibody (AAA127212). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel)/90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: rat C6 whole cell lysates, Lane 2: rat L6 whole cell lysates, Lane 3: mouse Neuro-2a whole cell lysates, Lane 4: mouse C2C12 whole cell lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-Osmr antigen affinity purified polyclonal antibody (#AAA127212) at 0.5ug/mL overnight at 4 degree C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for Osmr at approximately 110,180 kDa. The expected band size for Osmr is at 110 kDa.

ADDITIONAL INFORMATION

Related Product Information for anti-Osmr antibody	The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Product Categories/Family for anti-Osmr antibody	Primary Antibodies; Signal Transduction; Signaling Pathway; Small G Proteins
References	1. Arita, K., South, A. P., Hans-Filho, G., Sakuma, T. H., Lai-Cheong, J., Clements, S., Odashiro, M., Odashiro, D. N., Hans-Neto, G., Hans, N. R., Holder, M. V., Bhogal, B. S., Hartshorne, S. T., Akiyama, M., Shimizu, H., McGrath, J. A. Oncostatin M receptor-beta mutations underlie familial primary localized cutaneous amyloidosis. Am. J. Hum. Genet. 82: 73-80, 2008.2. Dillon, S. R., Sprecher, C., Hammond, A., Bilsborough, J., Rosenfeld-Franklin, M., Presnell, S. R., Haugen, H. S., Maurer, M., Harder, B., Johnston, J., Bort, S., Mudri, S., and 18 others. Interleukin 31, a cytokine produced by activated T cells, induces dermatitis in mice. Nature Immun. 5: 752-760, 2004. Note: Erratum: Nature Immun. 6: 114 only, 2005.3. Jahani-Asl, A., Yin, H., Soleimani, V. D., Haque, T., Luchman, H. A., Chang, N. C., Sincennes, M.-C., Puram, S. V., Scott, A. M., Lorimer, I. A. J., Perkins, T. J., Ligon, K. L., Weiss, S., Rudnicki, M. A., Bonni, A. Control of glioblastoma tumorigenesis by feed-forward cytokine signaling. Nature Neurosci. 19: 798-806, 2016.

NCBI AND UNIPROT INFORMATION

NCBI GI #	162287202
NCBI GeneID	18414
NCBI Accession #	NP_035149.2
NCBI GenBank Nucleotide #	NM_011019.3
UniProt Accession #	O70458; O88821
Molecular Weight	110,229 Da

NCBI Official Full Name	oncostatin-M-specific receptor subunit beta
UniProt Protein Name	Oncostatin-M-specific receptor subunit beta
UniProt Gene Name	Osmr
UniProt Synonym Gene Names	Osmrb; IL-31 receptor subunit beta; IL-31R subunit beta; IL-31R-beta; IL-31RB
UniProt Entry Name	OSMR_MOUSE

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/osmr-antibody-flow-cytometry-western-blot-elisa/127212>