

Mouse anti-Human CD22 Antibody | anti-CD22 antibody

Anti-Hu CD22 FITC


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA128628	Mouse	Human	Flow Cytometry, Functional Assay

PRODUCT SYNONYMS


CD22; Anti-Hu CD22 FITC; CD22 molecule; SIGLEC2; SIGLEC-2; anti-CD22 antibody

PRODUCT SPECIFICATIONS

Host	Mouse
Reactivity	Human
Isotype	IgG1
Clone Number	IS7
Specificity	The antibody IS7 reacts with an extracellular epitope of CD22 (BL-CAM), a 130 kDa type I transmembrane glycoprotein (immunoglobulin superfamily) expressed in the cytoplasm of pro-B and pre-B lymphocytes, and on the surface of mature and activated B lymphocytes; it is lost on plasma cells, peripheral blood T lymphocytes, granulocytes and monocytes.
Purity/Purification	The purified antibody is conjugated with fluorescein isothiocyanate (FITC) under optimum conditions. The conjugate is purified by size-exclusion chromatography.
Form/Format	Stabilizing phosphate buffered saline (PBS) solution containing 15 mM sodium azide
Sequence Length	759
Applicable Applications for anti-CD22 antibody	FCM/FACS (Flow Cytometry)
Immunogen	human cell line Reh
Conjugation	FITC
Preparation and Storage	Store in the dark at 2-8 degree C. Do not freeze. Avoid prolonged exposure to light.



FCM/FACS (Flow Cytometry)
Flow cytometry surface staining pattern of human peripheral whole blood stained using anti-human CD22 (IS7) FITC antibody (20 ul reagent / 100 ul of peripheral whole blood).



FCM/FACS (Flow Cytometry)
Separation of human CD22 positive lymphocytes (red-filled) from neutrophil granulocytes (black-dashed) in flow cytometry analysis (surface staining) of human peripheral whole blood stained using anti-human CD22 (IS7) FITC antibody (20 ul reagent / 100 ul of peripheral whole blood).

ADDITIONAL INFORMATION

Related Product Information for anti-CD22 antibody	CD22, also known as Siglec-2 (sialic acid-binding immunoglobulin-like lectin-2) is a transmembrane glycoprotein binding alpha2,6-linked sialic acid-bearing ligands. Intracellular domain of CD22 recruits protein tyrosine phosphatase SHP-1 through the immunoreceptor tyrosine-based inhibitory motifs (ITIMs), thus setting a threshold for B cell receptor-mediated activation. CD22 also regulates B-cell response by involvement in controlling the CD19/CD21-Src-family protein tyrosine kinase amplification pathway and CD40 signaling. CD22 exhibits hallmarks of clathrin-mediated endocytic pathway.
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	297374825
NCBI GeneID	933
NCBI Accession #	NP_001172028.1
NCBI GenBank Nucleotide #	NP_001172028.1
UniProt Accession #	P20273
NCBI Official Full Name	B-cell receptor CD22 isoform 2
NCBI Official Synonym Full Names	CD22 molecule
NCBI Official Symbol	CD22
NCBI Official Synonym Symbols	SIGLEC2; SIGLEC-2
NCBI Protein Information	B-cell receptor CD22
UniProt Protein Name	B-cell receptor CD22
UniProt Gene Name	CD22
UniProt Synonym Gene Names	SIGLEC2; BL-CAM; Siglec-2
UniProt Entry Name	CD22_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.