

Mouse anti-Human CD20 Antibody | anti-CD20 antibody

Anti-Hu CD20 FITC


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA128673	Mouse	Human	Flow Cytometry, Functional Assay

PRODUCT SYNONYMS


CD20; Anti-Hu CD20 FITC; MS4A1; B1; S7; MS4A; Bp35; CVID5; LEU-16; anti-CD20 antibody

PRODUCT SPECIFICATIONS

Host	Mouse
Reactivity	Human
Isotype	IgG2a
Clone Number	LT20
Specificity	The antibody LT20 reacts with an extracellular epitope of CD20 (Bp35), a 33-37 kDa non-glycosylated membrane receptor with four transmembrane domains, expressed on B lymphocytes (it is lost on plasma cells), follicular dendritic cells, and at low levels on peripheral blood T lymphocytes.
Purity/Purification	The purified antibody is conjugated with fluorescein isothiocyanate (FITC) under optimum conditions. The conjugate is purified by size-exclusion chromatography.
Form/Format	Stabilizing phosphate buffered saline (PBS) solution containing 15 mM sodium azide
Sequence Length	297
Applicable Applications for anti-CD20 antibody	FCM/FACS (Flow Cytometry)
Immunogen	Normal human lymphocytes from lymph node.
Conjugation	FITC
Preparation and Storage	Store in the dark at 2-8 degree C. Do not freeze. Avoid prolonged exposure to light.



FCM/FACS (Flow Cytometry)
Flow cytometry surface staining pattern of human peripheral whole blood stained using anti-human CD20 (LT20) FITC antibody (20 ul reagent / 100 ul of peripheral whole blood).



FCM/FACS (Flow Cytometry)
Separation of human CD20 positive lymphocytes (red-filled) from neutrophil granulocytes (black-dashed) in flow cytometry analysis (surface staining) of human peripheral whole blood stained using anti-human CD20 (LT20) FITC antibody (20 ul reagent / 100 ul of peripheral whole blood).

ADDITIONAL INFORMATION

Related Product Information for anti-CD20 antibody	CD20 is a cell surface 33-37 (depending on the degree of phosphorylation) kDa non-glycosylated surface phosphoprotein expressed on mature and most malignant B cells, but not stem cells or plasma cells (low number of the CD20 has been also detected on a subpopulation of T lymphocytes and it can be expressed on follicular dendritic cells). Its expression on B cells is synchronous with the expression of surface IgM. CD20 regulates transmembrane calcium conductance (probably functioning as a component of store-operated calcium channel), cell cycle progression and B-cell proliferation. It is associated with lipid rafts, but the intensity of this association depends on extracellular triggering, employing CD20 conformational change and/or BCR (B cell antigen receptor) aggregation. After the receptor ligation, BCR and CD20 colocalize and then rapidly dissociate before BCR endocytosis, whereas CD20 remains at the cell surface. CD20 serves as a useful target for antibody-mediated therapeutic depletion of B cells, as it is expressed at high levels on most B-cell malignancies, but does not become internalized or shed from the plasma membrane following mAb treatment.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	115968
NCBI GeneID	931
UniProt Accession #	P11836
NCBI Official Full Name	B-lymphocyte antigen CD20
NCBI Official Synonym Full Names	membrane spanning 4-domains A1
NCBI Official Symbol	MS4A1
NCBI Official Synonym Symbols	B1; S7; Bp35; CD20; CVID5; MS4A2; LEU-16
NCBI Protein Information	B-lymphocyte antigen CD20
UniProt Protein Name	B-lymphocyte antigen CD20
UniProt Gene Name	MS4A1
UniProt Synonym Gene Names	CD20
UniProt Entry Name	CD20_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

