

Mouse anti-Human CD3 Monoclonal Antibody | anti-CD3E antibody

Anti-Hu CD3 Purified

CATALOG # AAA128978	HOST Mouse	REACTIVITY Human	APPLICATIONS Immunocytochemistry, Flow Cytometry, Functional Assay
-------------------------------	----------------------	----------------------------	--

PRODUCT SYNONYMS

CD3; Monoclonal Antibody; Anti-Hu CD3 Purified; CD3 antigen; epsilon polypeptide; CD3E; T3E; TCRE; anti-CD3E antibody

PRODUCT SPECIFICATIONS

Host	Mouse
Reactivity	Human
Clonality	Monoclonal
Isotype	IgG2b
Clone Number	TB3
Specificity	The mouse monoclonal antibody TB3 recognizes an extracellular epitope on CD3 antigen of the TCR/CD3 complex on mature human T cells. This antibody, although not presented to HLDA workshops, has superior binding than the workshop clone TB2.
Purity/Purification	Purified by Protein-A Affinity Chromatography
Form/Format	Phosphate buffered saline (PBS) solution with 15 mM sodium azide
Concentration	1 mg/ml (varies by lot)
Sequence Length	207
Applicable Applications for anti-CD3E antibody	ICC (Immunocytochemistry), FCM/FACS (Flow Cytometry)
Preparation and Storage	Store at 2-8 degree C.

ADDITIONAL INFORMATION

Related Product Information for anti-CD3E antibody	CD3 complex is crucial in transducing antigen-recognition signals into the cytoplasm of T cells and in regulating the cell surface expression of the TCR complex. T cell activation through the antigen receptor (TCR) involves the cytoplasmic tails of the CD3 subunits CD3 gamma, CD3 delta, CD3 epsilon and CD3 zeta. These CD3 subunits are structurally related members of the immunoglobulins super family encoded by closely linked genes on human chromosome 11. The CD3 components have long cytoplasmic tails that associate with cytoplasmic signal transduction molecules. This association is mediated at least in part by a double tyrosine-based motif present in a single copy in the CD3 subunits. CD3 may play a role in TCR-induced growth arrest, cell survival and proliferation. The CD3 antigen is present on 68-82% of normal peripheral blood lymphocytes, 65-85% of thymocytes and Purkinje cells in the cerebellum. It is never expressed on B or NK cells. Decreased percentages of T lymphocytes may be observed in some autoimmune diseases.
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	1345708
NCBI GeneID	916
UniProt Accession #	P07766

NCBI Official Full Name	T-cell surface glycoprotein CD3 epsilon chain
NCBI Official Synonym Full Names	CD3e molecule
NCBI Official Symbol	CD3E
NCBI Official Synonym Symbols	T3E; TCRE; IMD18
NCBI Protein Information	T-cell surface glycoprotein CD3 epsilon chain
UniProt Protein Name	T-cell surface glycoprotein CD3 epsilon chain
UniProt Gene Name	CD3E
UniProt Synonym Gene Names	T3E
UniProt Entry Name	CD3E_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/monoclonal/cd3-antibody-clone-tb3-immunocytochemistry-flow-cytometry-elisa/128978>