

ACVR2B recombinant protein

Human ACVR2B Protein, hFc Tag

CATALOG #
AAA129495

HOST
HEK293


PRODUCT SYNONYMS

ACVR2B; Human ACVR2B Protein; hFc Tag; HTX4; ACTRIIB; ActR-IIB; Recombinant human ACVR2B Protein with C-terminal human Fc tag; ACVR2B recombinant protein

PRODUCT SPECIFICATIONS


Host	HEK293
Purity/Purification	The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.
Form/Format	Lyophilized from sterile PBS, pH 7.4. Normally 5% – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Sequence	ACVR2B(Ser 19-Thr137) hFc(Glu99- Ala330)
Tag	C-Human Fc
Preparation and Storage	Store at -20 degree C to -80 degree C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80 degree C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.

IMAGES



ELISA

Figure 2. ELISA plate pre-coated by 2 ug/mL (100 uL/well) Human ACVR2B Protein, hFc Tag (AAA129495) can bind Anti-ACVR2B(bimagrumab biosimilar) mAb in a linear range of 0.64–80 ng/mL.



SDS-PAGE

Figure 1. Human ACVR2B Protein, hFc Tag on SDS-PAGE under reducing condition.

ADDITIONAL INFORMATION

Related Product Information for ACVR2B recombinant protein	Activins are dimeric growth and differentiation factors which belong to the transforming growth factor-beta (TGF-beta) superfamily of structurally related signaling proteins. Activins signal through a heteromeric complex of receptor serine kinases which include at least two type I (I and IB) and two type II (II and IIB) receptors. These receptors are all transmembrane proteins, composed of a ligand-binding extracellular domain with cysteine-rich region, a transmembrane domain, and a cytoplasmic domain with predicted serine/threonine specificity. Type I receptors are essential for signaling; and type II receptors are required for binding ligands and for expression of type I receptors. Type I and II receptors form a stable complex after ligand binding, resulting in phosphorylation of type I receptors by type II receptors. Type II receptors are considered to be constitutively active kinases. This gene encodes activin A type IIB receptor, which displays a 3- to 4-fold higher affinity for the ligand than activin A type II receptor. [provided by RefSeq, Jul 2008]
Product Categories/Family for ACVR2B recombinant protein	ECD Proteins

NCBI AND UNIPROT INFORMATION

NCBI GI #	116734708
NCBI GeneID	93
NCBI Accession #	NP_001097.2
NCBI GenBank Nucleotide #	NM_001106.3
UniProt Accession #	Q13705; Q4VAV0
Molecular Weight	– Da
NCBI Official Full Name	activin receptor type-2B
NCBI Official Synonym Full Names	activin A receptor type 2B
NCBI Official Symbol	ACVR2B
NCBI Official Synonym Symbols	HTX4; ACTRIIB; ActR-IIB
NCBI Protein Information	activin receptor type-2B
UniProt Protein Name	Activin receptor type-2B
UniProt Gene Name	ACVR2B
UniProt Synonym Gene Names	ACTR-IIB
UniProt Entry Name	AVR2B_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/acvr2b-recombinant-protein-hek293/129495>