

Mouse Monokine Induced By Interferon Gamma (Mlg) CLIA Kit | Mlg clia kit

Mouse Monokine Induced By Interferon Gamma (Mlg) CLIA Kit

CATALOG #
AAA129920

REACTIVITY
Mouse


PRODUCT SYNONYMS

Monokine Induced By Interferon Gamma (Mlg); Mouse Monokine Induced By Interferon Gamma (Mlg) CLIA Kit; CXCL9; CMK; Humig; MI-G; SCYB9; Crg-10; Chemokine (C-X-C motif) ligand 9; Small-inducible cytokine B9; Gamma-interferon-induced monokine; Mlg clia kit

PRODUCT SPECIFICATIONS

Reactivity	Mouse
Specificity	This assay has high sensitivity and excellent specificity for detection of Monokine Induced By Interferon Gamma (Mlg). No significant cross-reactivity or interference between Monokine Induced By Interferon Gamma (Mlg) and analogues was observed.
Sequence Length	125
Assay Type	Double-antibody Sandwich
Samples	Serum, Plasma and other biological fluids
Detection Range	4.12-3,000pg/mL
Sensitivity	1.96pg/mL
Application	Chemiluminescent immunoassay for Antigen Detection.
Intra-assay Precision (Precision within an assay)	3 samples with low, middle and high level Monokine Induced By Interferon Gamma (Mlg) were tested 20 times on one plate, respectively.
Inter-assay Precision (Precision between assays)	3 samples with low, middle and high level Monokine Induced By Interferon Gamma (Mlg) were tested on 3 different plates, 8 replicates in each plate.
CV(%) =	SD/meanX100
Intra-Assay	CV<10%
Inter-Assay	CV<12%
Preparation and Storage	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for Mig clia kit	The microplate provided in this kit has been pre-coated with an antibody specific to Monokine Induced By Interferon Gamma (Mig). Standards or samples are then added to the appropriate microplate wells with a biotin-conjugated antibody specific to Monokine Induced By Interferon Gamma (Mig). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. Then the mixture of substrate A and B is added to generate glow light emission kinetics. Upon plate development, the intensity of the emitted light is proportional to the Monokine Induced By Interferon Gamma (Mig) level in the sample or standard.;
Product Categories/Family for Mig clia kit	Cytokine; Infection immunity

NCBI AND UNIPROT INFORMATION

NCBI GI #	161579144
NCBI GeneID	513990
UniProt Accession #	A9QWP9
Molecular Weight	14,106 Da
NCBI Official Full Name	monokine induced by interferon gamma
NCBI Official Symbol	CXCL9
NCBI Protein Information	C-X-C motif chemokine 9
UniProt Protein Name	C-X-C motif chemokine 9
UniProt Gene Name	CXCL9
UniProt Entry Name	CXCL9_BOVIN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/mouse-clia-kits/monokine-induced-by-interferon-gamma-mig/129920>