

Rat Calreticulin (CRT) CLIA Kit | CRT clia kit

Rat Calreticulin (CRT) CLIA Kit

CATALOG #
AAA130057

REACTIVITY
Rat


PRODUCT SYNONYMS

Calreticulin (CRT); Rat Calreticulin (CRT) CLIA Kit; CALR; RO; SSA; CC1qR; ERp60; HACBP; grp60; CRTC; CRP55; Calregulin; Sicca Syndrome Antigen A; Autoantigen Ro; Endoplasmic reticulum resident protein 60; CRT clia kit

PRODUCT SPECIFICATIONS

Reactivity	Rat
Specificity	This assay has high sensitivity and excellent specificity for detection of Calreticulin (CRT). No significant cross-reactivity or interference between Calreticulin (CRT) and analogues was observed.
Sequence Length	417
Assay Type	Double-antibody Sandwich
Samples	Serum, Plasma and other biological fluids
Detection Range	2.74-2,000pg/mL
Sensitivity	1.21pg/mL
Application	Chemiluminescent immunoassay for Antigen Detection.
Intra-assay Precision (Precision within an assay)	3 samples with low, middle and high level Calreticulin (CRT) were tested 20 times on one plate, respectively.
Inter-assay Precision (Precision between assays)	3 samples with low, middle and high level Calreticulin (CRT) were tested on 3 different plates, 8 replicates in each plate.
CV(%) =	SD/meanX100
Intra-Assay	CV<10%
Inter-Assay	CV<12%
Preparation and Storage	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for CRT clia kit	The microplate provided in this kit has been pre-coated with an antibody specific to Calreticulin (CRT). Standards or samples are then added to the appropriate microplate wells with a biotin-conjugated antibody specific to Calreticulin (CRT). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. Then the mixture of substrate A and B is added to generate glow light emission kinetics. Upon plate development, the intensity of the emitted light is proportional to the Calreticulin (CRT) level in the sample or standard.;
Product Categories/Family for CRT clia kit	Signal transduction; Metabolic pathway; Bone metabolism

NCBI AND UNIPROT INFORMATION

NCBI GI #	4757900
NCBI GeneID	811
NCBI Accession #	NP_004334.1
NCBI GenBank Nucleotide #	NM_004343.3
UniProt Accession #	P27797; Q6IAT4; Q9UDG2
Molecular Weight	48,142 Da
NCBI Official Full Name	calreticulin
NCBI Official Synonym Full Names	calreticulin
NCBI Official Symbol	CALR
NCBI Official Synonym Symbols	RO; CRT; SSA; cC1qR; HEL-S-99n
NCBI Protein Information	calreticulin
UniProt Protein Name	Calreticulin
UniProt Gene Name	CALR
UniProt Synonym Gene Names	CRTC; ERp60
UniProt Entry Name	CALR_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/rat-clia-kits/calreticulin-crt/130057>