

Human Ceruloplasmin (CP) CLIA Kit | CP clia kit

Human Ceruloplasmin (CP) CLIA Kit

CATALOG #
AAA130097

REACTIVITY
Human


PRODUCT SYNONYMS

Ceruloplasmin (CP); Human Ceruloplasmin (CP) CLIA Kit; CP2; CER; Caeruloplasmin; Ferroxidase; Oxygen Oxidoreductase; CP clia kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of Ceruloplasmin (CP). No significant cross-reactivity or interference between Ceruloplasmin (CP) and analogues was observed.
Sequence Length	1065
Assay Type	Double-antibody Sandwich
Samples	Serum, Plasma, Tissue homogenates, Cell culture supernates and other biological fluids
Detection Range	1.37-1,000pg/mL
Sensitivity	0.58pg/mL
Application	Chemiluminescent immunoassay for Antigen Detection.
Intra-assay Precision (Precision within an assay)	3 samples with low, middle and high level Ceruloplasmin (CP) were tested 20 times on one plate, respectively.
Inter-assay Precision (Precision between assays)	3 samples with low, middle and high level Ceruloplasmin (CP) were tested on 3 different plates, 8 replicates in each plate.
CV(%) =	SD/meanX100
Intra-Assay	CV<10%
Inter-Assay	CV<12%
Preparation and Storage	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for CP clia kit	The microplate provided in this kit has been pre-coated with an antibody specific to Ceruloplasmin (CP). Standards or samples are then added to the appropriate microplate wells with a biotin-conjugated antibody specific to Ceruloplasmin (CP). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. Then the mixture of substrate A and B is added to generate glow light emission kinetics. Upon plate development, the intensity of the emitted light is proportional to the Ceruloplasmin (CP) level in the sample or standard.;
Product Categories/Family for CP clia kit	Metabolic pathway ; Infection immunity

NCBI AND UNIPROT INFORMATION

NCBI GI #	4557485
NCBI GeneID	1356
NCBI Accession #	NP_000087.1
NCBI GenBank Nucleotide #	NM_000096.3
UniProt Accession #	P00450 ; Q14063 ; Q2PP18 ; Q9UKS4
Molecular Weight	122,205 Da
NCBI Official Full Name	ceruloplasmin
NCBI Official Synonym Full Names	ceruloplasmin (ferroxidase)
NCBI Official Symbol	CP
NCBI Official Synonym Symbols	CP-2
NCBI Protein Information	ceruloplasmin
UniProt Protein Name	Ceruloplasmin
UniProt Gene Name	CP
UniProt Entry Name	CERU_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.