

Rabbit Glycated Hemoglobin A1c (HbA1c) Polyclonal Antibody | anti-HbA1c antibody

Polyclonal Antibody to Glycated Hemoglobin A1c (HbA1c)

CATALOG # AAA131659	HOST Rabbit	REACTIVITY Homo sapiens (Human)	APPLICATIONS Immunohistochemistry, Western Blot
-------------------------------	-----------------------	---	---

PRODUCT SYNONYMS

Glycated Hemoglobin A1c (HbA1c); Polyclonal Antibody; Polyclonal Antibody to Glycated Hemoglobin A1c (HbA1c); anti-HbA1c antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Homo sapiens (Human)
Clonality	Polyclonal
Purity/Purification	Antigen-specific affinity chromatography followed by Protein A affinity chromatography
Form/Format	Supplied as solution form in 0.01M PBS, pH7.4, containing 0.05% Proclin-300, 50% glycerol.
Concentration	0.5mg/mL (varies by lot)
Applicable Applications for anti-HbA1c antibody	IHC (Immunohistochemistry), WB (Western Blot)
Source	Polyclonal antibody preparation
Traits	Liquid
Cross Reactivity	Mouse; Rat
Immunogen	Native HbA1c .
Preparation and Storage	Storage: Avoid repeated freeze/thaw cycles. Store at 4 degree C for frequent use. Aliquot and store at -20 degree C for 24 months. Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37 degree C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

IMAGES



IHC (Immunohistochemistry)

DAB staining on IHC-P; Samples: Human Colon Tissue; Primary Ab: 20ug/ml Rabbit Anti-Human HbA1c Antibody; Second Ab: 2ug/mL HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody



IHC (Immunohistochemistry)


DAB staining on IHC-P; Samples: Human Spleen Tissue; Primary Ab: 20ug/ml Rabbit Anti-Human HbA1c Antibody; Second Ab: 2ug/mL HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody

**IHC (Immunohistochemistry)**


DAB staining on IHC-P; Samples: Human Spinal cord Tissue; Primary Ab: 20ug/ml Rabbit Anti-Human HbA1C Antibody Second Ab: 2ug/mL HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody

**IHC (Immunohistochemistry)**

DAB staining on IHC-P; Samples: Human Kidney Tissue; Primary Ab: 20ug/ml Rabbit Anti-Human HbA1C Antibody Second Ab: 2ug/mL HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody

**IHC (Immunohistochemistry)**

DAB staining on IHC-P; Samples: Human Cardiac Muscle Tissue; Primary Ab: 20ug/ml Rabbit Anti-Human HbA1C Antibody Second Ab: 2ug/mL HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody

**WB (Western Blot)**

Western Blot: Lane 1: Rat Blood Cells Lane 2: Mouse Blood Cells

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/glycated-hemoglobin-a1c-hba1c-antibody-immunohistochemistry-western-blot-elisa/131659>