

Rabbit Procollagen C Proteinase Enhancer 2 (PCPE2) Polyclonal Antibody | anti-PCPE2 antibody

Polyclonal Antibody to Procollagen C Proteinase Enhancer 2 (PCPE2)

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA132007	Rabbit	Homo sapiens (Human)	Immunohistochemistry, Immunofluorescence, Immunocytochemistry


PRODUCT SYNONYMS

Procollagen C Proteinase Enhancer 2 (PCPE2); Polyclonal Antibody; Polyclonal Antibody to Procollagen C Proteinase Enhancer 2 (PCPE2); anti-PCPE2 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Homo sapiens (Human)
Clonality	Polyclonal
Purity/Purification	Antigen-specific affinity chromatography followed by Protein A affinity chromatography
Form/Format	Supplied as solution form in 0.01M PBS, pH7.4, containing 0.05% Proclin-300, 50% glycerol.
Concentration	0.5mg/mL (varies by lot)
Applicable Applications for anti-PCPE2 antibody	IHC (Immunohistochemistry), IF (Immunofluorescence), ICC (Immunocytochemistry)
Source	Polyclonal antibody preparation
Traits	Liquid
Cross Reactivity	Porcine
Immunogen	Recombinant PCPE2 (Cys297~Cys415 (Accession # Q9UKZ9)) expressed in E.coli()
Preparation and Storage	Storage: Avoid repeated freeze/thaw cycles. Store at 4 degree C for frequent use. Aliquot and store at -20 degree C for 24 months. Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37 degree C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

IMAGES



IF (Immunofluorescence)

FITC staining on IF;Sample: Human U2OS cell;Primary Ab: 30ug/ml Rabbit Anti-Human PCPE2 AntibodySecond Ab: 2ug/ml FITC-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody



IHC (Immunohistochemistry)


DAB staining on IHC-P;Sample: Porcine Testis Tissue;Primary Ab: 20ug/ml Rabbit Anti-Human PCPE2 AntibodySecond Ab: 2ug/mL HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody

**IHC (Immunohistochemistry)**

DAB staining on IHC-P; Sample: Porcine Lymph node Tissue; Primary Ab: 20ug/ml Rabbit Anti-Human PCPE2 Antibody; Second Ab: 2ug/mL HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody

**IHC (Immunohistochemistry)**

DAB staining on IHC-P; Sample: Porcine Cerebrum Tissue; Primary Ab: 20ug/ml Rabbit Anti-Human PCPE2 Antibody; Second Ab: 2ug/mL HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody

**IHC (Immunohistochemistry)**

DAB staining on IHC-P; Samples: Human Cerebrum Tissue; Primary Ab: 20ug/ml Rabbit Anti-Human PCPE2 Antibody; Second Ab: 2ug/mL HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody

NCBI AND UNIPROT INFORMATION

NCBI GI #	7019483
NCBI GeneID	26577
NCBI Accession #	NP_037495.1
NCBI GenBank Nucleotide #	NM_013363.3
UniProt Accession #	Q9UKZ9; Q9BRH3; B2RCH9; D3DNG4
NCBI Official Full Name	procollagen C-endopeptidase enhancer 2
NCBI Official Synonym Full Names	procollagen C-endopeptidase enhancer 2
NCBI Official Symbol	PCOLCE2
NCBI Official Synonym Symbols	PCPE2
NCBI Protein Information	procollagen C-endopeptidase enhancer 2
UniProt Protein Name	Procollagen C-endopeptidase enhancer 2
UniProt Gene Name	PCOLCE2
UniProt Synonym Gene Names	PCPE2; PCPE-2; Procollagen C-proteinase enhancer 2

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/procollagen-c-proteinase-enhancer-2-pcpe2-antibody-immunohistochemistry-immunofluorescence-immunocytochemistry-elisa/132007>