

Rabbit Protein C Receptor, Endothelial (PROCR) Polyclonal Antibody | anti-PROCR antibody

Polyclonal Antibody to Protein C Receptor, Endothelial (PROCR)

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA133620	Rabbit	Rat, Human, Mouse	Immunoprecipitation, Immunocytochemistry, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

Protein C Receptor; Endothelial (PROCR); Polyclonal Antibody; Polyclonal Antibody to Protein C Receptor; anti-PROCR antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Rat, Human, Mouse
Clonality	Polyclonal
Specificity	The antibody is a Rabbit polyclonal antibody raised against PROCR. It has been selected for its ability to recognize PROCR in immunohistochemical staining and western blotting.
Purity/Purification	Antigen-specific Affinity Chromatography
Form/Format	Supplied as solution form in 0.01M PBS, pH7.4, containing 0.05% Proclin-300, 50% glycerol
Concentration	1mg/ml (varies by lot)
Applicable Applications for anti-PROCR antibody	IP (Immunoprecipitation), ICC (Immunocytochemistry), IHC (Immunohistochemistry), WB (Western Blot)
Source	Polyclonal antibody preparation
Traits	Liquid
Immunogen	Recombinant PROCR (Glu59~Gly217) expressed in E.coli
Cross Reactivity	Rat
Conjugated Antibody	The APC conjugated antibody version of this item is also available as
Preparation and Storage	Storage: Avoid repeated freeze/thaw cycles. Store at 4 degree C for frequent use. Aliquot and store at -20 degree C for 12 months. Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37 degree C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.


IMAGES




IHC (Immunohistochemistry)
DAB staining on IHC-P; Samples: Rat Testis Tissue.




IHC (Immunohistochemistry)
DAB staining on IHC-P; Samples: Rat Kidney Tissue.



IHC (Immunohistochemistry)
DAB staining on IHC-P; Samples: Rat Intestine Tissue.



IHC (Immunohistochemistry)
DAB staining on IHC-P; Samples: Rat Pancreas Tissue



WB (Western Blot)
Western Blot: Sample: Recombinant PROCR, Rat.

NCBI AND UNIPROT INFORMATION

NCBI GI #	71043878
NCBI GeneID	362248
NCBI Accession #	NP_001020904.1
NCBI GenBank Nucleotide #	NM_001025733.2
UniProt Accession #	Q4V8I1
Molecular Weight	26,839 Da
NCBI Official Full Name	endothelial protein C receptor
NCBI Official Synonym Full Names	protein C receptor

NCBI Official Symbol	Procr
NCBI Protein Information	endothelial protein C receptor
UniProt Protein Name	Endothelial protein C receptor
UniProt Gene Name	Procr
UniProt Synonym Gene Names	APC receptor

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/protein-c-receptor-endothelial-procr-antibody-immunoprecipitation-immunocytochemistry-immunohistochemistry-western-blot-elisa/133620>