

Cluster Of Differentiation 8a (CD8a) Recombinant Protein | CD8a recombinant protein

Recombinant Cluster Of Differentiation 8a (CD8a)

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA134287	E Coli	Human	Immunoprecipitation, Immunoprecipitation, ELISA, Western Blot, SDS-Page

PRODUCT SYNONYMS

Cluster Of Differentiation 8a (CD8a); Recombinant Cluster Of Differentiation 8a (CD8a); CD8a recombinant protein

PRODUCT SPECIFICATIONS

Host	E Coli
Reactivity	Human
Purity/Purification	>92%
Form/Format	20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% sarcosyl, 5%Trehalose and Proclin300
Concentration	200ug/mL (varies by lot)
Applicable Applications for CD8a recombinant protein	IP (Coimmunoprecipitation), IP (Immunoprecipitation), ELISA, WB (Western Blot), SDS-PAGE
Source	Prokaryotic expression.
Residues	Ser22~Val198
Tags	N-terminal His-Tag
Subcellular Location	Cell membrane; Single-pass type I membrane protein. Secreted.
Traits	Freeze-dried powder
Predicted isoelectric point	8.8
Predicted Molecular Mass	20.6kDa
Accurate Molecular Mass	24&28kDa as determined by SDS-PAGE reducing conditions
Phenomenon explanation	The possible reasons that the actual band size differs from the predicted are as follows: 1. Splice variants: Alternative splicing may create different sized proteins from the same gene. 2. Relative charge: The composition of amino acids may affects the charge of the protein. 3. Post-translational modification: Phosphorylation, glycosylation, methylation etc. 4. Post-translation cleavage: Many proteins are synthesized as pro-proteins, and then cleaved to give the active form. 5. Polymerization of the target protein: Dimerization, multimerization etc.
Usage	Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.
Preparation and Storage	Storage: Avoid repeated freeze/thaw cycles. Store at 2-8 degree C for one month. Aliquot and store at -80 degree C for 12 months. Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37 degree C for 48h, and no obvious degradation and precipitation were observed.The loss rate is less than 5% within the expiration date under appropriate storage condition.

IMAGES

Application Data



Sequence



SDS-PAGE



NCBI AND UNIPROT INFORMATION

NCBI GI #	225007536
NCBI GeneID	925
NCBI Accession #	NP_001139345.1
NCBI GenBank Nucleotide #	NM_001145873.1
UniProt Accession #	P01732; Q13970; Q4ZG17; D6W5M8
NCBI Official Full Name	T-cell surface glycoprotein CD8 alpha chain isoform 1
NCBI Official Synonym Full Names	CD8a molecule
NCBI Official Symbol	CD8A
NCBI Official Synonym Symbols	CD8; MAL; p32; Leu2
NCBI Protein Information	T-cell surface glycoprotein CD8 alpha chain; T8 T-cell antigen; T cell co-receptor; OKT8 T-cell antigen; T-cell antigen Leu2; Leu2 T-lymphocyte antigen; CD8 antigen, alpha polypeptide (p32); T-lymphocyte differentiation antigen T8/Leu-2
UniProt Protein Name	T-cell surface glycoprotein CD8 alpha chain
UniProt Gene Name	CD8A
UniProt Synonym Gene Names	MAL
UniProt Entry Name	CD8A_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.