

Human Tubulin Beta (TUBb) ELISA Kit | TUBb elisa kit

Human Tubulin Beta (TUBb) ELISA Kit

CATALOG #
AAA136691

REACTIVITY
Human


PRODUCT SYNONYMS

Tubulin Beta (TUBb); Human Tubulin Beta (TUBb) ELISA Kit; TUBb elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of Tubulin Beta (TUBb) No significant cross-reactivity or interference between Tubulin Beta (TUBb) and analogues was observed.
Sequence Length	439
Samples	Serum, plasma, tissue homogenates, Cell lysates and other biological fluids
Assay Type	Sandwich
Detection Range	31.25-2000pg/mL The standard curve concentrations used for the ELISA's were 2000pg/mL, 1000pg/mL, 500pg/mL, 250pg/mL, 125pg/mL, 62.5pg/mL, 31.25pg/mL
Sensitivity	Typically less than 12.7pg/mL.
Organism Species	Homo sapiens (Human)
ELISA Type	Sandwich
Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level Tubulin Beta (TUBb) were tested 20 times on one plate, respectively. Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level Tubulin Beta (TUBb) were tested on 3 different plates, 8 replicates in each plate. $CV(\%) = SD/mean \times 100$ Intra-Assay: $CV < 10\%$ Inter-Assay: $CV < 12\%$
Preparation and Storage	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for TUBb elisa kit	The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate provided in this kit has been pre-coated with an antibody specific to Tubulin Beta (TUBb). Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody specific to Tubulin Beta (TUBb). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain Tubulin Beta (TUBb), biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of Tubulin Beta (TUBb) in the samples is then determined by comparing the O.D. of the samples to the standard curve.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	358342735
NCBI Official Full Name	tubulin beta

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/human-elisa-kits/tubulin-beta-tubb/136691>