

Goat TNFRSF1A/TNFR1 Polyclonal Antibody | anti-TNFRSF1A antibody

Goat anti-TNFRSF1A/TNFR1 Antibody

CATALOG # AAA13672	HOST Goat	REACTIVITY Tested: Human; Expected from sequence similarity: Human, Rat, Pig	APPLICATIONS ELISA, Western Blot
------------------------------	---------------------	--	--


PRODUCT SYNONYMS

TNFRSF1A/TNFR1; Polyclonal Antibody; Goat anti-TNFRSF1A/TNFR1 Antibody; TNFRSF1A; tumor necrosis factor receptor superfamily; member 1A; CD120a; FPF; MGC19588; TBP1; TNF-R; TNF-R-I; TNF-R55; TNFAR; TNFR1; TNFR55; TNFR60; p55; p55-R; p60; TNF-R1; TNF-R1; TNFR-I; tumor necrosis factor binding protein 1; tumor necrosis factor re; CD120a antibody; FPF antibody; MGC19588 antibody; p55 antibody; p55-R antibody; p60 antibody; TBP1 antibody; TNFAR antibody; TNF-R antibody; TNFR1 antibody; TNF-R1 antibody; TNFR55 antibody; TNF-R55 antibody; TNFR60 antibody; TNFR-I antibody; TNF-R1 antibody; TNF-R-I antibody; tumor necrosis factor binding protein 1 antibody; tumor necrosis factor receptor 1A isoform beta antibody; tumor necrosis factor receptor superfamily member 1A antibody; member 1A antibody; tumor necrosis factor receptor type 1 antibody; tumor necrosis factor-alpha receptor antibody; TNFRSF1A antibody; TNFRSF1A / TNFR1; anti-TNFRSF1A antibody


PRODUCT SPECIFICATIONS

Host	Goat
Reactivity	Tested: Human; Expected from sequence similarity: Human, Rat, Pig
Clonality	Polyclonal
Purity/Purification	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Form/Format	Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.
Concentration	100ug specific antibody in 200ul (varies by lot)
Sequence	SDHEIDRLEL QNGR
Sequence Length	455
Applicable Applications for anti-TNFRSF1A antibody	ELISA (Peptide ELISA), WB (Western Blot)
Application Notes	Peptide ELISA: Antibody detection limit dilution 1: 32000. Western Blot: Approx 52kDa band observed in Human Lung lysates and in lysates of cell lines A549, HepG2 and K562 (calculated MW of 50.5kDa according to NP_001056.1). Recommended concentration: 0.3-1ug/ml.
Immunogen	Peptide with sequence C-SDHEIDRLELQNGR, from the internal region of the protein sequence according to NP_001056.1.
Epitope	Internal region
Preparation and Storage	Aliquot and store at -20 degree C. Minimize freezing and thawing.

IMAGES



WB (Western Blot)
0.5ug/ml staining of A549 (A), HepG2 (B) and K562 (C) lysates (35ug protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.



WB (Western Blot)
0.3ug/ml staining of Human Lung lysate (35ug protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

NCBI AND UNIPROT INFORMATION

NCBI GI #	4507575
NCBI GeneID	7132
NCBI Accession #	NP_001056.1
NCBI GenBank Nucleotide #	NM_001065.3
NCBI Official Full Name	tumor necrosis factor receptor superfamily member 1A
NCBI Official Synonym Full Names	tumor necrosis factor receptor superfamily member 1A
NCBI Official Symbol	TNFRSF1A
NCBI Official Synonym Symbols	FPF; MS5; p55; p60; TBP1; TNF-R; TNFAR; TNFR1; p55-R; CD120a; TNFR55; TNFR60; TNF-R-I; TNF-R55; TNFR1-d2
NCBI Protein Information	tumor necrosis factor receptor superfamily member 1A
UniProt Protein Name	Tumor necrosis factor receptor superfamily member 1A
UniProt Gene Name	TNFRSF1A
UniProt Synonym Gene Names	TNFAR; TNFR1; TNF-R1; TNF-RI; TNFR-I; TBPI
UniProt Entry Name	TNR1A_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/tnfrsf1a-tnfr1-antibody-peptide-elisa-western-blot-elisa/13672>