

Cytochrome C, Somatic (CYCS) ELISA Kit | CYCS elisa kit

Cytochrome C, Somatic (CYCS) ELISA Kit

CATALOG #
AAA136838


PRODUCT SYNONYMS

Cytochrome C; Somatic (CYCS); Somatic (CYCS) ELISA Kit; CYCS elisa kit

PRODUCT SPECIFICATIONS

Specificity	This assay has high sensitivity and excellent specificity for detection of Cytochrome C, Somatic (CYCS). No significant cross-reactivity or interference between Cytochrome C, Somatic (CYCS) and analogues was observed.
Sequence Length	105
Samples	Serum, plasma, tissue homogenates and other biological fluids
Assay Type	Sandwich
Detection Range	1.563-100ng/mL The standard curve concentrations used for the ELISA's were 100ng/mL, 50ng/mL, 25ng/mL, 12.5ng/mL, 6.25ng/mL, 3.125ng/mL, 1.563ng/mL
Sensitivity	Typically less than 0.57ng/mL.
Organism Species	Mus musculus (Mouse), Rattus norvegicus (Rat)
ELISA Type	Sandwich
Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level Cytochrome C, Somatic (CYCS) were tested 20 times on one plate, respectively. Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level Cytochrome C, Somatic (CYCS) were tested on 3 different plates, 8 replicates in each plate. $CV(\%) = SD/mean \times 100$ Intra-Assay: $CV < 10\%$ Inter-Assay: $CV < 12\%$
Preparation and Storage	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for CYCS elisa kit	The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate provided in this kit has been pre-coated with an antibody specific to Cytochrome C, Somatic (CYCS). Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody specific to Cytochrome C, Somatic (CYCS). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain Cytochrome C, Somatic (CYCS), biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of Cytochrome C, Somatic (CYCS) in the samples is then determined by comparing the O.D. of the samples to the standard curve.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	45501024
NCBI GeneID	54205
UniProt Accession #	P99999; P00001; Q6NUR2; Q6NX69; Q96BV4; A4D166; B2R411
Molecular Weight	11,749 Da
NCBI Official Full Name	Cytochrome c, somatic
NCBI Official Synonym Full Names	cytochrome c, somatic
NCBI Official Symbol	CYCS
NCBI Official Synonym Symbols	CYC; HCS; THC4
NCBI Protein Information	cytochrome c
UniProt Protein Name	Cytochrome c
UniProt Gene Name	CYCS
UniProt Synonym Gene Names	CYC
UniProt Entry Name	CYC_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.