

Rat Tenascin C (TNC) ELISA Kit | TNC elisa kit

Rat Tenascin C (TNC) ELISA Kit

CATALOG #
AAA136917

REACTIVITY
Rat


PRODUCT SYNONYMS

Tenascin C (TNC); Rat Tenascin C (TNC) ELISA Kit; TN-C; HXB; TN; GMEM; JI; Myotendinous antigen; Neuronectin; GP 150-225; Cytotactin; Hexabrachion; Glioma-associated-extracellular matrix antigen; TNC elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Rat
Specificity	This assay has high sensitivity and excellent specificity for detection of Tenascin C (TNC). No significant cross-reactivity or interference between Tenascin C (TNC) and analogues was observed.
Sequence Length	91
Assay Type	Double-antibody Sandwich
Samples	Serum, Plasma, Tissue homogenates, Cell lysates, Cell culture supernates and other biological fluids
Detection Range	0.312-20ng/mL
Sensitivity	0.131ng/mL
Application	Enzyme-linked immunosorbent assay for Antigen Detection.
Intra-assay Precision (Precision within an assay)	3 samples with low, middle and high level Tenascin C (TNC) were tested 20 times on one plate, respectively.
Inter-assay Precision (Precision between assays)	3 samples with low, middle and high level Tenascin C (TNC) were tested on 3 different plates, 8 replicates in each plate.
CV(%) =	SD/meanX100
Intra-Assay	CV<10%
Inter-Assay	CV<12%
Preparation and Storage	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for TNC elisa kit	The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate provided in this kit has been pre-coated with an antibody specific to Tenascin C (TNC). Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody specific to Tenascin C (TNC). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain Tenascin C (TNC), biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of Tenascin C (TNC) in the samples is then determined by comparing the O.D. of the samples to the standard curve.
Product Categories/Family for TNC elisa kit	Metabolic pathway; Neuroscience; Developmental science; Nutrition metabolism; Bone metabolism

NCBI AND UNIPROT INFORMATION

NCBI GI #	1842130
Molecular Weight	9,764 Da
NCBI Official Full Name	tenascin-C, partial
UniProt Protein Name	Tenascin-C
UniProt Entry Name	Q99857_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/rat-elisa-kits/tenascin-c-tnc/136917>