Human Chemokine C-C-Motif Ligand 14 (CCL14) ELISA Kit | CCL14 elisa kit

Human Chemokine C-C-Motif Ligand 14 (CCL14) ELISA Kit

CATALOG #	REACTIVITY
AAA137338	**Human**


PRODUCT SYNONYMS

Chemokine C-C-Motif Ligand 14 (CCL14); Human Chemokine C-C-Motif Ligand 14 (CCL14) ELISA Kit; SCYA14; HCC-1; HCC-3; NCC-2; SCYL2; CKb1; MCIF; Small Inducible Cytokine Subfamily A(Cys-Cys)Member 14; CCL14 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of Chemokine C-C-Motif Ligand 14 (CCL14). No significant cross-reactivity or interference between Chemokine C-C-Motif Ligand 14 (CCL14) and analogues was observed.
Sequence Length	28
Assay Type	Double-antibody Sandwich
Samples	Serum, Plasma, Tissue homogenates, Cell lysates, Cell culture supernates and other biological fluids
Detection Range	15.6-1000pg/mL
Sensitivity	6.1pg/mL
Application	Enzyme-linked immunosorbent assay for Antigen Detection.
Intra-assay Precision (Precision within an assay)	3 samples with low, middle and high level Chemokine C-C-Motif Ligand 14 (CCL14) were tested 20 times on one plate, respectively.
Inter-assay Precision (Precision between assays)	3 samples with low, middle and high level Chemokine C-C-Motif Ligand 14 (CCL14) were tested on 3 different plates, 8 replicates in each plate.
CV(%) =	SD/meanX100
Intra-Assay	CV<10%
Inter-Assay	CV<12%
Preparation and Storage	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for CCL14 elisa kit	The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate provided in this kit has been pre-coated with an antibody specific to Chemokine C-C-Motif Ligand 14 (CCL14). Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody specific to Chemokine C-C-Motif Ligand 14 (CCL14). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain Chemokine C-C-Motif Ligand 14 (CCL14), biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of Chemokine C-C-Motif Ligand 14 (CCL14) in the samples is then determined by comparing the O.D. of the samples to the standard curve.
Product Categories/Family for CCL14 elisa kit	Cytokine; Tumor immunity; Infection immunity; Immune molecule

NCBI AND UNIPROT INFORMATION

NCBI GI #	805489819
Molecular Weight	3,362 Da
NCBI Official Full Name	chemokine (C-C motif) ligand 14, partial
UniProt Protein Name	Chemokine (C-C motif) ligand 14
UniProt Gene Name	**CCL14**
UniProt Entry Name	A0A0E3SUY6_OCHDA

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: https://stage.aaabiotech.com/human-elisa-kits/chemokine-c-c-motif-ligand-14-ccl14/137338