

Human Fc Fragment Of IgG Receptor Transporter Alpha (FcγRT) ELISA Kit | FcγRT elisa kit

Human Fc Fragment Of IgG Receptor Transporter Alpha (FcγRT) ELISA Kit

CATALOG #
AAA137593

REACTIVITY
Human


PRODUCT SYNONYMS

Fc Fragment Of IgG Receptor Transporter Alpha (FcγRT); Human Fc Fragment Of IgG Receptor Transporter Alpha (FcγRT) ELISA Kit; FCRN; Neonatal Fc receptor; IgG receptor FcRn large subunit p51; FcγRT elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of FcγRI. No significant cross-reactivity or interference between FcγRI and analogues was observed.
Sequence Length	365
Assay Type	Double-antibody Sandwich
Samples	Tissue homogenates, Cell lysates and other biological fluids
Detection Range	0.156-10ng/mL
Sensitivity	0.057ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level FcγRI were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level FcγRI were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for FcgRT elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of FcgRI in human serum, plasma, tissue homogenates and other biological fluids. Principle of the Assay: The microplate provided in this kit has been pre-coated with an antibody specific to FcgRI. Standards or samples are then added to the appropriate microplate wells with a biotin-conjugated antibody specific to FcgRI. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain FcgRI, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of FcgRI in the samples is then determined by comparing the O.D. of the samples to the standard curve.
Product Categories/Family for FcgRT elisa kit	CD & Adhesion molecule

NCBI AND UNIPROT INFORMATION

NCBI GI #	14250561
NCBI GeneID	2217
UniProt Accession #	P55899; Q5HYM5; Q9HBV7; Q9NZ19
Molecular Weight	39,743 Da
NCBI Official Full Name	Fc fragment of IgG, receptor, transporter, alpha
NCBI Official Synonym Full Names	Fc fragment of IgG, receptor, transporter, alpha
NCBI Official Symbol	FCGRT
NCBI Official Synonym Symbols	FCRN; alpha-chain
NCBI Protein Information	IgG receptor FcRn large subunit p51
UniProt Protein Name	IgG receptor FcRn large subunit p51
UniProt Gene Name	FCGRT
UniProt Synonym Gene Names	FCRN; FcRn
UniProt Entry Name	FCGRN_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/human-elisa-kits/fc-fragment-of-igg-receptor-transporter-alpha-fcgrt/137593>