

Mouse Monokine Induced By Interferon Gamma (Mlg) ELISA Kit | Mlg elisa kit

Mouse Monokine Induced By Interferon Gamma (Mlg) ELISA Kit

CATALOG #
AAA138147

REACTIVITY
Mouse


PRODUCT SYNONYMS

Monokine Induced By Interferon Gamma (Mlg); Mouse Monokine Induced By Interferon Gamma (Mlg) ELISA Kit; CXCL9; CMK; Humig; MI-G; SCYB9; Crg-10; Chemokine (C-X-C motif) ligand 9; Small-inducible cytokine B9; Gamma-interferon-induced monokine; Mlg elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Mouse
Specificity	This assay has high sensitivity and excellent specificity for detection of Monokine Induced By Interferon Gamma (Mlg). No significant cross-reactivity or interference between Monokine Induced By Interferon Gamma (Mlg) and analogues was observed.
Sequence Length	125
Assay Type	Double-antibody Sandwich
Samples	Serum, Plasma, Tissue homogenates, Cell lysates, Cell culture supernates and other biological fluids
Detection Range	15.6-1,000pg/mL
Sensitivity	5.8pg/mL
Application	Enzyme-linked immunosorbent assay for Antigen Detection.
Intra-assay Precision (Precision within an assay)	3 samples with low, middle and high level Monokine Induced By Interferon Gamma (Mlg) were tested 20 times on one plate, respectively.
Inter-assay Precision (Precision between assays)	3 samples with low, middle and high level Monokine Induced By Interferon Gamma (Mlg) were tested on 3 different plates, 8 replicates in each plate.
CV(%) =	SD/meanX100
Intra-Assay	CV<10%
Inter-Assay	CV<12%
Preparation and Storage	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for MIg elisa kit	The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate provided in this kit has been pre-coated with an antibody specific to Monokine Induced By Interferon Gamma (MIg). Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody specific to Monokine Induced By Interferon Gamma (MIg). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain Monokine Induced By Interferon Gamma (MIg), biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of Monokine Induced By Interferon Gamma (MIg) in the samples is then determined by comparing the O.D. of the samples to the standard curve.
Product Categories/Family for MIg elisa kit	Cytokine; Infection immunity

NCBI AND UNIPROT INFORMATION

NCBI GI #	161579144
NCBI GeneID	513990
UniProt Accession #	A9QWP9
Molecular Weight	14,106 Da
NCBI Official Full Name	monokine induced by interferon gamma
NCBI Official Symbol	CXCL9
NCBI Protein Information	C-X-C motif chemokine 9
UniProt Protein Name	C-X-C motif chemokine 9
UniProt Gene Name	CXCL9
UniProt Entry Name	CXCL9_BOVIN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/mouse-elisa-kits/monokine-induced-by-interferon-gamma-mig/138147>