

Human Endocrine Gland Derived Vascular Endothelial Growth Factor (EG-VEGF) ELISA Kit

| EG-VEGF elisa kit

Human Endocrine Gland Derived Vascular Endothelial Growth Factor (EG-VEGF) ELISA Kit

CATALOG #
AAA138162

REACTIVITY
Human


PRODUCT SYNONYMS

Endocrine Gland Derived Vascular Endothelial Growth Factor (EG-VEGF); Human Endocrine Gland Derived Vascular Endothelial Growth Factor (EG-VEGF) ELISA Kit; PK1; EGVEGF; PRK1; PROK1; Prokineticin 1; Black Mamba Toxin-Related Protein; Mambakine; EG-VEGF elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of Endocrine Gland Derived Vascular Endothelial Growth Factor (EG-VEGF). No significant cross-reactivity or interference between Endocrine Gland Derived Vascular Endothelial Growth Factor (EG-VEGF) and analogues was observed.
Sequence Length	105
Assay Type	Double-antibody Sandwich
Samples	Serum, Plasma, Tissue homogenates, Cell lysates, Cell culture supernates and other biological fluids
Detection Range	15.6-1,000pg/mL
Sensitivity	6.5pg/mL
Application	Enzyme-linked immunosorbent assay for Antigen Detection.
Intra-assay Precision (Precision within an assay)	3 samples with low, middle and high level Endocrine Gland Derived Vascular Endothelial Growth Factor (EG-VEGF) were tested 20 times on one plate, respectively.
Inter-assay Precision (Precision between assays)	3 samples with low, middle and high level Endocrine Gland Derived Vascular Endothelial Growth Factor (EG-VEGF) were tested on 3 different plates, 8 replicates in each plate.
CV(%) =	SD/meanX100
Intra-Assay	CV<10%
Inter-Assay	CV<12%
Preparation and Storage	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for EG-VEGF elisa kit	The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate provided in this kit has been pre-coated with an antibody specific to Endocrine Gland Derived Vascular Endothelial Growth Factor (EG-VEGF). Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody specific to Endocrine Gland Derived Vascular Endothelial Growth Factor (EG-VEGF). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain Endocrine Gland Derived Vascular Endothelial Growth Factor (EG-VEGF), biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of Endocrine Gland Derived Vascular Endothelial Growth Factor (EG-VEGF) in the samples is then determined by comparing the O.D. of the samples to the standard curve.
Product Categories/Family for EG-VEGF elisa kit	Cytokine; Tumor immunity

NCBI AND UNIPROT INFORMATION

NCBI GI #	37182488
NCBI GeneID	84432
UniProt Accession #	P58294; Q5VWD4; Q8TC69
Molecular Weight	11,715 Da
NCBI Official Full Name	EG-VEGF
NCBI Official Synonym Full Names	prokineticin 1
NCBI Official Symbol	PROK1
NCBI Official Synonym Symbols	PK1; PRK1; EGVEGF
NCBI Protein Information	prokineticin-1
UniProt Protein Name	Prokineticin-1
UniProt Gene Name	PROK1
UniProt Synonym Gene Names	EG-VEGF
UniProt Entry Name	PROK1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.