

Human Chemokine C-X-C-Motif Receptor 4 (CXCR4) ELISA Kit | CXCR4 elisa kit

Human Chemokine C-X-C-Motif Receptor 4 (CXCR4) ELISA Kit

CATALOG #
AAA138198

REACTIVITY
Human


PRODUCT SYNONYMS

Chemokine C-X-C-Motif Receptor 4 (CXCR4); Human Chemokine C-X-C-Motif Receptor 4 (CXCR4) ELISA Kit; CD184; Fusin; NPYR; HSY3RR; LAP3; LCR1; LESTR; NPY3R; NPYRL; WHIM; Leukocyte-derived seven transmembrane domain receptor; Lipopolysaccharide-associated protein 3; CXCR4 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of Chemokine C-X-C-Motif Receptor 4 (CXCR4). No significant cross-reactivity or interference between Chemokine C-X-C-Motif Receptor 4 (CXCR4) and analogues was observed.
Sequence Length	356
Assay Type	Double-antibody Sandwich
Samples	Tissue homogenates, Cell lysates and other biological fluids
Detection Range	0.156-10ng/mL
Sensitivity	0.056ng/mL
Application	Enzyme-linked immunosorbent assay for Antigen Detection.
Intra-assay Precision (Precision within an assay)	3 samples with low, middle and high level Chemokine C-X-C-Motif Receptor 4 (CXCR4) were tested 20 times on one plate, respectively.
Inter-assay Precision (Precision between assays)	3 samples with low, middle and high level Chemokine C-X-C-Motif Receptor 4 (CXCR4) were tested on 3 different plates, 8 replicates in each plate.
CV(%) =	SD/meanX100
Intra-Assay	CV<10%
Inter-Assay	CV<12%
Preparation and Storage	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for CXCR4 elisa kit	The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate provided in this kit has been pre-coated with an antibody specific to Chemokine C-X-C-Motif Receptor 4 (CXCR4). Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody specific to Chemokine C-X-C-Motif Receptor 4 (CXCR4). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain Chemokine C-X-C-Motif Receptor 4 (CXCR4), biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of Chemokine C-X-C-Motif Receptor 4 (CXCR4) in the samples is then determined by comparing the O.D. of the samples to the standard curve.
Product Categories/Family for CXCR4 elisa kit	Signal transduction; CD & Adhesion molecule; Tumor immunity; Infection immunity

NCBI AND UNIPROT INFORMATION

NCBI GI #	29825383
NCBI GeneID	7852
UniProt Accession #	P61073; O60835; P30991; P56438; Q53S69; Q9BXA0; Q9UKN2; B2R5N0
Molecular Weight	40,221 Da
NCBI Official Full Name	chemokine (C-X-C motif) receptor 4
NCBI Official Synonym Full Names	chemokine (C-X-C motif) receptor 4
NCBI Official Symbol	CXCR4
NCBI Official Synonym Symbols	FB22; HM89; LAP3; LCR1; NPYR; WHIM; CD184; LAP-3; LESTR; NPY3R; NPYRL; WHIMS; HSY3RR; NPY3R; D2S201E
NCBI Protein Information	C-X-C chemokine receptor type 4
UniProt Protein Name	C-X-C chemokine receptor type 4
UniProt Gene Name	CXCR4
UniProt Synonym Gene Names	CXC-R4; CXCR-4; LESTR; LAP-3; LPS-associated protein 3; SDF-1 receptor
UniProt Entry Name	CXCR4_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.