

Human BMP And Activin Membrane Bound Inhibitor Homolog (BAMBI) ELISA Kit | BAMBI elisa kit

Human BMP And Activin Membrane Bound Inhibitor Homolog (BAMBI) ELISA Kit

CATALOG # AAA138584	REACTIVITY Human
-------------------------------	----------------------------


PRODUCT SYNONYMS

BMP And Activin Membrane Bound Inhibitor Homolog (BAMBI); Human BMP And Activin Membrane Bound Inhibitor Homolog (BAMBI) ELISA Kit; NMA; Non-metastatic gene A protein; Putative transmembrane protein NMA; BAMBI elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of BMP And Activin Membrane Bound Inhibitor Homolog (BAMBI). No significant cross-reactivity or interference between BMP And Activin Membrane Bound Inhibitor Homolog (BAMBI) and analogues was observed.
Sequence Length	260
Assay Type	Double-antibody Sandwich
Samples	Tissue homogenates, Cell lysates and other biological fluids
Detection Range	0.156-10ng/mL
Sensitivity	0.059ng/mL
Application	Enzyme-linked immunosorbent assay for Antigen Detection.
Intra-assay Precision (Precision within an assay)	3 samples with low, middle and high level BMP And Activin Membrane Bound Inhibitor Homolog (BAMBI) were tested 20 times on one plate, respectively.
Inter-assay Precision (Precision between assays)	3 samples with low, middle and high level BMP And Activin Membrane Bound Inhibitor Homolog (BAMBI) were tested on 3 different plates, 8 replicates in each plate.
CV(%) =	SD/meanX100
Intra-Assay	CV<10%
Inter-Assay	CV<12%
Preparation and Storage	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for BAMBI elisa kit	The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate provided in this kit has been pre-coated with an antibody specific to BMP And Activin Membrane Bound Inhibitor Homolog (BAMBI). Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody specific to BMP And Activin Membrane Bound Inhibitor Homolog (BAMBI). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain BMP And Activin Membrane Bound Inhibitor Homolog (BAMBI), biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of BMP And Activin Membrane Bound Inhibitor Homolog (BAMBI) in the samples is then determined by comparing the O.D. of the samples to the standard curve.
Product Categories/Family for BAMBI elisa kit	Immune molecule

NCBI AND UNIPROT INFORMATION

NCBI GI #	6912534
NCBI GeneID	25805
NCBI Accession #	NP_036474.1
NCBI GenBank Nucleotide #	NM_012342.2
UniProt Accession #	Q13145
Molecular Weight	29,108 Da
NCBI Official Full Name	BMP and activin membrane-bound inhibitor homolog
NCBI Official Synonym Full Names	BMP and activin membrane-bound inhibitor
NCBI Official Symbol	BAMBI
NCBI Official Synonym Symbols	NMA
NCBI Protein Information	BMP and activin membrane-bound inhibitor homolog
UniProt Protein Name	BMP and activin membrane-bound inhibitor homolog
UniProt Gene Name	BAMBI
UniProt Synonym Gene Names	NMA
UniProt Entry Name	BAMBI_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.