

Human Procollagen III (PCIII) ELISA Kit | PCIII elisa kit

Human Procollagen III (PCIII) ELISA Kit

CATALOG #
AAA138745

REACTIVITY
Human


PRODUCT SYNONYMS

Procollagen III (PCIII); Human Procollagen III (PCIII) ELISA Kit; PCOL-III; PC-III; PC3; Type III Procollagen; PCIII elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of Procollagen III (PCIII). No significant cross-reactivity or interference between Procollagen III (PCIII) and analogues was observed.
Sequence Length	68
Assay Type	Double-antibody Sandwich
Samples	Serum, Plasma, Tissue homogenates and other biological fluids
Detection Range	0.156-10ng/mL
Sensitivity	0.059ng/mL
Application	Enzyme-linked immunosorbent assay for Antigen Detection.
Intra-assay Precision (Precision within an assay)	3 samples with low, middle and high level Procollagen III (PCIII) were tested 20 times on one plate, respectively.
Inter-assay Precision (Precision between assays)	3 samples with low, middle and high level Procollagen III (PCIII) were tested on 3 different plates, 8 replicates in each plate.
CV(%) =	SD/meanX100
Intra-Assay	CV<10%
Inter-Assay	CV<12%
Preparation and Storage	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for PCIII elisa kit	The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate provided in this kit has been pre-coated with an antibody specific to Procollagen III (PCIII). Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody specific to Procollagen III (PCIII). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain Procollagen III (PCIII), biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of Procollagen III (PCIII) in the samples is then determined by comparing the O.D. of the samples to the standard curve.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	119587121
NCBI GeneID	5119
UniProt Accession #	Q9HD42; Q14468; Q15779; Q96G31; A2RU09
Molecular Weight	7,363 Da
NCBI Official Full Name	procollagen (type III) N-endopeptidase, isoform CRA_a, partial
NCBI Official Synonym Full Names	charged multivesicular body protein 1A
NCBI Official Symbol	CHMP1A
NCBI Official Synonym Symbols	PCH8; CHMP1; PRSM1; PCOLN3; VPS46A; VPS46-1
NCBI Protein Information	charged multivesicular body protein 1a
UniProt Protein Name	Charged multivesicular body protein 1a
UniProt Gene Name	CHMP1A
UniProt Synonym Gene Names	; CHMP1a; Vps46-1; hVps46-1
UniProt Entry Name	CHM1A_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.