

Rat N-Acetyltransferase 2 (NAT2) ELISA Kit | NAT2 elisa kit

Rat N-Acetyltransferase 2 (NAT2) ELISA Kit

CATALOG #
AAA138810

REACTIVITY
Rat


PRODUCT SYNONYMS

N-Acetyltransferase 2 (NAT2); Rat N-Acetyltransferase 2 (NAT2) ELISA Kit; AAC2; PNAT; Arylamine N-Acetyltransferase; Arylamide acetylase 2; Polymorphic arylamine N-acetyltransferase; NAT2 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Rat
Specificity	This assay has high sensitivity and excellent specificity for detection of N-Acetyltransferase 2 (NAT2). No significant cross-reactivity or interference between N-Acetyltransferase 2 (NAT2) and analogues was observed.
Sequence Length	290
Assay Type	Double-antibody Sandwich
Samples	Tissue homogenates and other biological fluids
Detection Range	0.625-40ng/mL
Sensitivity	0.218ng/mL
Application	Enzyme-linked immunosorbent assay for Antigen Detection.
Intra-assay Precision (Precision within an assay)	3 samples with low, middle and high level N-Acetyltransferase 2 (NAT2) were tested 20 times on one plate, respectively.
Inter-assay Precision (Precision between assays)	3 samples with low, middle and high level N-Acetyltransferase 2 (NAT2) were tested on 3 different plates, 8 replicates in each plate.
CV(%) =	SD/meanX100
Intra-Assay	CV<10%
Inter-Assay	CV<12%
Preparation and Storage	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for NAT2 elisa kit	The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate provided in this kit has been pre-coated with an antibody specific to N-Acetyltransferase 2 (NAT2). Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody specific to N-Acetyltransferase 2 (NAT2). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain N-Acetyltransferase 2 (NAT2), biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of N-Acetyltransferase 2 (NAT2) in the samples is then determined by comparing the O.D. of the samples to the standard curve.
Product Categories/Family for NAT2 elisa kit	Enzyme & Kinase

NCBI AND UNIPROT INFORMATION

NCBI GI #	29423790
NCBI GeneID	10
UniProt Accession #	P11245; O43637; O60654; O60655; Q13146; Q16697; Q2MLE4; Q2MLF5; Q2MLG8; Q2MLJ6; Q2MLK4; Q2MLK6
Molecular Weight	33,542 Da
NCBI Official Full Name	N-acetyltransferase 2
NCBI Official Synonym Full Names	N-acetyltransferase 2 (arylamine N-acetyltransferase)
NCBI Official Symbol	NAT2
NCBI Official Synonym Symbols	AAC2; PNAT; NAT-2
NCBI Protein Information	arylamine N-acetyltransferase 2
UniProt Protein Name	Arylamine N-acetyltransferase 2
UniProt Gene Name	NAT2
UniProt Synonym Gene Names	AAC2; NAT-2; PNAT
UniProt Entry Name	ARY2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.