

Human Intercellular Adhesion Molecule 4 (ICAM4) ELISA Kit | ICAM4 elisa kit

Human Intercellular Adhesion Molecule 4 (ICAM4) ELISA Kit

CATALOG #
AAA139514

REACTIVITY
Human


PRODUCT SYNONYMS

Intercellular Adhesion Molecule 4 (ICAM4); Human Intercellular Adhesion Molecule 4 (ICAM4) ELISA Kit; CD242; LW; Landsteiner-Weiner Blood System; Landsteiner-Wiener blood group glycoprotein; ICAM4 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of ICAM1. No significant cross-reactivity or interference between ICAM1 and analogues was observed.
Sequence Length	272
Assay Type	Double-antibody Sandwich
Samples	Tissue homogenates, Cell lysates and other biological fluids
Detection Range	0.156-10ng/mL
Sensitivity	0.063ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level ICAM1 were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level ICAM1 were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for ICAM4 elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of ICAM1 in mouse serum, plasma, tissue homogenates, cell lysates, cell culture supernates and other biological fluids. Principle of the Assay: The microplate provided in this kit has been pre-coated with an antibody specific to ICAM1. Standards or samples are then added to the appropriate microplate wells with a biotin-conjugated antibody specific to ICAM1. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain ICAM1, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of ICAM1 in the samples is then determined by comparing the O.D. of the samples to the standard curve.
Product Categories/Family for ICAM4 elisa kit	CD & Adhesion molecule; Tumor immunity; Infection immunity; Hematology

NCBI AND UNIPROT INFORMATION

NCBI GI #	4504561
NCBI GeneID	3386
NCBI Accession #	NP_001535.1
NCBI GenBank Nucleotide #	NM_001544.4
UniProt Accession #	Q14773; Q14771; Q14772; Q16375; Q9BWR0; A0M8X2
Molecular Weight	29,649 Da
NCBI Official Full Name	intercellular adhesion molecule 4 isoform 1
UniProt Protein Name	Intercellular adhesion molecule 4
UniProt Gene Name	ICAM4
UniProt Synonym Gene Names	LW; ICAM-4; LW blood group protein
UniProt Entry Name	ICAM4_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/human-elisa-kits/intercellular-adhesion-molecule-4-icam4/139514>