

Rabbit Platelet Factor 4 (PF4) ELISA Kit | PF4 elisa kit

Rabbit Platelet Factor 4 (PF4) ELISA Kit

CATALOG #
AAA139727

REACTIVITY
Rabbit


PRODUCT SYNONYMS

Platelet Factor 4 (PF4); Rabbit Platelet Factor 4 (PF4) ELISA Kit; CXCL4; SCYB4; Chemokine C-X-C-Motif Ligand 4; Oncostatin-A; Iroplact; PF4 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Rabbit
Specificity	This assay has high sensitivity and excellent specificity for detection of PF4. No significant cross-reactivity or interference between PF4 and analogues was observed.
Sequence Length	101
Assay Type	Quantitative Sandwich
Samples	Rabbit platelet-poor plasma
Detection Range	78-5,000pg/mL
Sensitivity	< 30pg/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level PF4 were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level PF4 were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for PF4 elisa kit	Principle of the Assay: The microplate provided in this kit has been pre-coated with an antibody specific to PF4. Standards or samples are then added to the appropriate microplate wells with a biotin-conjugated antibody specific to PF4. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain PF4, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of PF4 in the samples is then determined by comparing the O.D. of the samples to the standard curve. Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of PF4 in rabbit platelet-poor plasma.
Product Categories/Family for PF4 elisa kit	Cytokine; Hematology

NCBI AND UNIPROT INFORMATION

NCBI GI #	4505733
NCBI GeneID	5196
NCBI Accession #	NP_002610.1
NCBI GenBank Nucleotide #	NM_002619.3
UniProt Accession #	P02776; Q53X61; Q9UC64; Q9UC65
Molecular Weight	10,845 Da
NCBI Official Full Name	platelet factor 4
NCBI Official Synonym Full Names	platelet factor 4
NCBI Official Symbol	PF4
NCBI Official Synonym Symbols	PF-4; CXCL4; SCYB4
NCBI Protein Information	platelet factor 4
UniProt Protein Name	Platelet factor 4
UniProt Gene Name	PF4
UniProt Synonym Gene Names	CXCL4; SCYB4; PF-4
UniProt Entry Name	PLF4_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.