

Human Toll Like Receptor 4 (TLR4) ELISA Kit | TLR4 elisa kit

Human Toll Like Receptor 4 (TLR4) ELISA Kit

CATALOG #
AAA13996

REACTIVITY
Human


PRODUCT SYNONYMS

Toll Like Receptor 4 (TLR4); Human Toll Like Receptor 4 (TLR4) ELISA Kit; CD284; TOLL; HToll; TLR4 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of TLR4. No significant cross-reactivity or interference between TLR4 and analogues was observed.
Sequence Length	799
Samples	Human tissue homogenates, cell lysates, cell culture supernates or other biological fluids
Assay Type	Quantitative Sandwich
Detection Range	0.156-10ng/mL
Sensitivity	< 0.057ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level TLR4 were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level TLR4 were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for TLR4 elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for the in vitro quantitative measurement of TLR4 in human tissue homogenates, cell lysates, cell culture supernates or other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to TLR4. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to TLR4. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain TLR4, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of TLR4 in the samples is then determined by comparing the O.D. of the samples to the standard curve.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	4507533
NCBI GeneID	7099
NCBI Accession #	NP_003257.1
NCBI GenBank Nucleotide #	NM_003266.3
UniProt Accession #	O00206; Q5VZI8; Q5VZI9; Q9UK78; Q9UM57; A8K1Y8; A9XLP9; A9XLQ0; A9XLQ1; B4E194; D1CS52; D1CS53
Molecular Weight	73,301 Da
NCBI Official Full Name	toll-like receptor 4 isoform C
NCBI Official Synonym Full Names	toll like receptor 4
NCBI Official Symbol	TLR4
NCBI Official Synonym Symbols	TOLL; CD284; TLR-4; ARMD10
NCBI Protein Information	toll-like receptor 4
UniProt Protein Name	Toll-like receptor 4
UniProt Gene Name	TLR4

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.