

Human Programmed Cell Death Protein 1 Ligand 2 (PDCD1LG2) ELISA Kit | PDCD1LG2 elisa kit

Human Programmed Cell Death Protein 1 Ligand 2 (PDCD1LG2) ELISA Kit

CATALOG #
AAA140290

REACTIVITY
Human


PRODUCT SYNONYMS

Programmed Cell Death Protein 1 Ligand 2 (PDCD1LG2); Human Programmed Cell Death Protein 1 Ligand 2 (PDCD1LG2) ELISA Kit; CD273; B7DC; Btdc; PD-L2; PDCD1-L2; PDL2; PDCD1LG2 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of Programmed Cell Death Protein 1 Ligand 2 (PDL2). No significant cross-reactivity or interference between Programmed Cell Death Protein 1 Ligand 2 (PDL2) and analogues was observed.
Sequence Length	182
Assay Type	Double-antibody Sandwich
Samples	Tissue homogenates, Cell lysates, Cell culture supernates and other biological fluids
Detection Range	0.156-10ng/mL
Sensitivity	0.061ng/mL
Application	Enzyme-linked immunosorbent assay for Antigen Detection.
Intra-assay Precision (Precision within an assay)	3 samples with low, middle and high level Programmed Cell Death Protein 1 Ligand 2 (PDL2) were tested 20 times on one plate, respectively.
Inter-assay Precision (Precision between assays)	3 samples with low, middle and high level Programmed Cell Death Protein 1 Ligand 2 (PDL2) were tested on 3 different plates, 8 replicates in each plate.
CV(%) =	SD/meanX100
Intra-Assay	CV<10%
Inter-Assay	CV<12%
Preparation and Storage	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for PDCD1LG2 elisa kit	The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate provided in this kit has been pre-coated with an antibody specific to Programmed Cell Death Protein 1 Ligand 2 (PDL2). Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody specific to Programmed Cell Death Protein 1 Ligand 2 (PDL2). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain Programmed Cell Death Protein 1 Ligand 2 (PDL2), biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of Programmed Cell Death Protein 1 Ligand 2 (PDL2) in the samples is then determined by comparing the O.D. of the samples to the standard curve.
Product Categories/Family for PDCD1LG2 elisa kit	Signal transduction; Tumor immunity; Infection immunity; Immune molecule; Autoimmunity

NCBI AND UNIPROT INFORMATION

NCBI GI #	190014605
NCBI GeneID	80380
NCBI Accession #	NP_079515.2
NCBI GenBank Nucleotide #	NM_025239.3
UniProt Accession #	Q9BQ51; Q14CN8; Q5T7Z6; Q6JXL8; Q6JXL9
Molecular Weight	20,745 Da
NCBI Official Full Name	programmed cell death 1 ligand 2
NCBI Official Synonym Full Names	programmed cell death 1 ligand 2
NCBI Official Symbol	PDCD1LG2
NCBI Official Synonym Symbols	B7DC; Btdc; PDL2; CD273; PD-L2; PDCD1L2; bA574F11.2
NCBI Protein Information	programmed cell death 1 ligand 2
UniProt Protein Name	Programmed cell death 1 ligand 2
UniProt Gene Name	PDCD1LG2
UniProt Synonym Gene Names	B7DC; CD273; PDCD1L2; PDL2; PD-1 ligand 2; PD-L2; PDCD1 ligand 2; Programmed death ligand 2; B7-DC
UniProt Entry Name	PD1L2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.