

Mouse Intercellular Adhesion Molecule 4 (ICAM4) ELISA Kit | ICAM4 elisa kit

Mouse Intercellular Adhesion Molecule 4 (ICAM4) ELISA Kit


CATALOG # AAA141184	REACTIVITY Mouse
-------------------------------	----------------------------

PRODUCT SYNONYMS

Intercellular Adhesion Molecule 4 (ICAM4); Mouse Intercellular Adhesion Molecule 4 (ICAM4) ELISA Kit; CD242; LW; Landsteiner-Weiner Blood System; Landsteiner-Wiener blood group glycoprotein; ICAM4 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Mouse
Specificity	This assay has high sensitivity and excellent specificity for detection of Intercellular Adhesion Molecule 4 (ICAM4). No significant cross-reactivity or interference between Intercellular Adhesion Molecule 4 (ICAM4) and analogues was observed.
Sequence Length	271
Assay Type	Double-antibody Sandwich
Samples	Tissue homogenates, Cell lysates and other biological fluids
Detection Range	0.156-10ng/mL
Sensitivity	0.059ng/mL
Application	Enzyme-linked immunosorbent assay for Antigen Detection.
Intra-assay Precision (Precision within an assay)	3 samples with low, middle and high level Intercellular Adhesion Molecule 4 (ICAM4) were tested 20 times on one plate, respectively.
Inter-assay Precision (Precision between assays)	3 samples with low, middle and high level Intercellular Adhesion Molecule 4 (ICAM4) were tested on 3 different plates, 8 replicates in each plate.
CV(%) =	SD/meanX100
Intra-Assay	CV<10%
Inter-Assay	CV<12%
Preparation and Storage	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for ICAM4 elisa kit	The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate provided in this kit has been pre-coated with an antibody specific to Intercellular Adhesion Molecule 4 (ICAM4). Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody specific to Intercellular Adhesion Molecule 4 (ICAM4). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain Intercellular Adhesion Molecule 4 (ICAM4), biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of Intercellular Adhesion Molecule 4 (ICAM4) in the samples is then determined by comparing the O.D. of the samples to the standard curve.
Product Categories/Family for ICAM4 elisa kit	CD & Adhesion molecule; Tumor immunity; Infection immunity; Hematology

NCBI AND UNIPROT INFORMATION

NCBI GI #	4504561
NCBI GeneID	3386
NCBI Accession #	NP_001535.1
NCBI GenBank Nucleotide #	NM_001544.4
Molecular Weight	29,649 Da
NCBI Official Full Name	intercellular adhesion molecule 4 isoform 1
NCBI Official Synonym Full Names	intercellular adhesion molecule 4 (Landsteiner-Wiener blood group)
NCBI Official Symbol	ICAM4
NCBI Official Synonym Symbols	LW; CD242
NCBI Protein Information	intercellular adhesion molecule 4
UniProt Protein Name	Intercellular adhesion molecule 4
UniProt Gene Name	ICAM4
UniProt Synonym Gene Names	LW; ICAM-4; LW blood group protein
UniProt Entry Name	ICAM4_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.