

Rabbit Histone H4 Polyclonal Antibody | anti-H4 antibody

Polyclonal Antibody to Histone H4 (H4)


CATALOG # AAA142844	HOST Rabbit	APPLICATIONS Western Blot
-------------------------------	-----------------------	-------------------------------------

PRODUCT SYNONYMS


Histone H4; Polyclonal Antibody; Polyclonal Antibody to Histone H4 (H4); anti-H4 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Clonality	Polyclonal
Purity/Purification	Antigen-specific affinity chromatography followed by Protein A affinity chromatography
Form/Format	Liquid; Supplied as solution form in 0.01M PBS, pH7.4, containing 0.05% Proclin-300, 50% glycerol.
Concentration	0.5mg/mL (varies by lot)
Applicable Applications for anti-H4 antibody	WB (Western Blot)
Source	Polyclonal antibody preparation
Cross Reactivity	Human;Mouse;Porcine
Immunogen	Recombinant Histone H4 (Ser2~Gly103) expressed in E.coli
Preparation and Storage	Storage: Avoid repeated freeze/thaw cycles. Store at 4°C for frequent use. Aliquot and store at -20°C for 24 months. Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.



WB (Western Blot)
Western Blot: Sample: Recombinant H4, Human.



WB (Western Blot)
Western Blot: Sample: Lane1: Porcine Spleen Tissue; Lane2: Mouse Thymus Tissue; Lane3: Mouse Placenta Tissue; Lane4: Human Hela Cells.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.