

Rabbit Mediator Complex Subunit 1 Polyclonal Antibody | anti-MED1 antibody

Mediator Complex Subunit 1 (MED1) Polyclonal Antibody

CATALOG # AAA144689	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS ELISA, Immunohistochemistry, Immunocytochemistry, Western Blot
-------------------------------	-----------------------	--	---


PRODUCT SYNONYMS

Mediator Complex Subunit 1; Polyclonal Antibody; Mediator Complex Subunit 1 (MED1) Polyclonal Antibody; PPARBP; ARC205; PBP; CRSP1; CRSP200; PPARGBP; RB18A; TRAP220; TRIP2; Peroxisome proliferator-activated receptor-binding protein; Thyroid receptor-interacting protein 2; anti-MED1 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Specificity	The antibody is a rabbit polyclonal antibody raised against MED1. It has been selected for its ability to recognize MED1 in immunohistochemical staining and western blotting.
Purity/Purification	Affinity Chromatography
Form/Format	Supplied as solution form in PBS, pH7.4 containing 0.02% NaN ₃ , 50% glycerol.
Concentration	500ug/mL (varies by lot)
Sequence Length	1581
Applicable Applications for anti-MED1 antibody	ELISA, IHC (Immunohistochemistry), ICC (Immunocytochemistry), WB (Western Blot)
Immunoglobulin Type	IgG
Conjugate	No Conjugate
Cross Reactivity	Human
Organism Species	Human
Fragment	MED1 (Met1~Thr212)
Quality Control	Content: The quality control contains recombinant MED1 (Met1~Thr212) disposed in loading buffer. Usage: 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate. 5uL per well when used in enhanced chemilumescient (ECL). Note: The quality control is specifically manufactured as the positive control. Not used for other purposes. Loading Buffer: 100mM Tris(pH8.8), 2% SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, NaN ₃ 0.02%.
Conjugated Antibody	The APC conjugated antibody version of this item is also available as
Preparation and Storage	Store at 4 degree C for frequent use. Stored at -20 degree C to -80 degree C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.

IMAGES



IHC (Immunohiostchemistry)
DAB staining on fromalin fixed paraffin- embedded rectum tissue



WB (Western Blot)
Western Blot: Sample: Recombinant MED1, Human.

NCBI AND UNIPROT INFORMATION

NCBI GI #	28559039
NCBI GeneID	5469
NCBI Accession #	NP_004765.2
NCBI GenBank Nucleotide #	NM_004774.3
UniProt Accession #	Q15648; O43810; O75447; Q6P9H7; Q6PK58; Q9HD39; A2RRQ6
Molecular Weight	61,563 Da
NCBI Official Full Name	mediator of RNA polymerase II transcription subunit 1
NCBI Official Synonym Full Names	mediator complex subunit 1
NCBI Official Symbol	MED1
NCBI Official Synonym Symbols	PBP; CRSP1; RB18A; TRIP2; PPARBP; CRSP200; DRIP205; DRIP230; PPARGBP; TRAP220
NCBI Protein Information	mediator of RNA polymerase II transcription subunit 1
UniProt Protein Name	Mediator of RNA polymerase II transcription subunit 1
UniProt Gene Name	MED1
UniProt Synonym Gene Names	ARC205; CRSP1; CRSP200; DRIP205; DRIP230; PBP; PPARBP; PPARGBP; RB18A; TRAP220; TRIP2; ARC205; PBP; PPAR-binding protein; Trap220; TR-interacting protein 2; TRIP-2

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/mediator-complex-subunit-1-antibody-immunohistochemistry-immunocytochemistry-western-blot-elisa/144689>