

Rabbit anti-Human Signaling Lymphocytic Activation Molecule Family Polyclonal Antibody | anti-SLAMF2 antibody

Signaling Lymphocytic Activation Molecule Family, Member 2 (SLAMF2) Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA144743	Rabbit	Human	ELISA, Immunohistochemistry, Immunocytochemistry, Western Blot

PRODUCT SYNONYMS

Signaling Lymphocytic Activation Molecule Family; Polyclonal Antibody; Member 2 (SLAMF2) Polyclonal Antibody; CD48; BLAST; BLAST1; MEM-102; BCM1; HCD48; MCD48; B-lymphocyte activation marker BLAST-1; BCM1 surface antigen; Leukocyte antigen MEM-102; SLAM family member 2; anti-SLAMF2 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Specificity	The antibody is a rabbit polyclonal antibody raised against SLAMF2. It has been selected for its ability to recognize SLAMF2 in immunohistochemical staining and western blotting.
Purity/Purification	Affinity Chromatography
Form/Format	Supplied as solution form in PBS, pH7.4 containing 0.02% NaN ₃ , 50% glycerol.
Concentration	0.25mg/ml (varies by lot)
Sequence Length	335
Applicable Applications for anti-SLAMF2 antibody	ELISA, IHC (Immunohistochemistry), ICC (Immunocytochemistry), WB (Western Blot)
Immunoglobulin Type	IgG
Conjugate	No Conjugate
Cross Reactivity	Human
Organism Species	Human
Fragment	SLAMF2 (Gln27~Thr243)
Quality Control	Content: The quality control contains recombinant SLAMF2 (Gln27~Thr243) disposed in loading buffer. Usage: 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate. 5uL per well when used in enhanced chemilumescence (ECL). Note: The quality control is specifically manufactured as the positive control. Not used for other purposes. Loading Buffer: 100mM Tris(pH8.8), 2% SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, NaN ₃ 0.02%.
Conjugated Antibody	The APC conjugated antibody version of this item is also available as
Preparation and Storage	Store at 4 degree C for frequent use. Stored at -20 degree C to -80 degree C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.

IMAGES



IHC (Immunohistochemistry)
DAB staining on formalin fixed paraffin-embedded kidney tissue



WB (Western Blot)
Western Blot: Sample: Recombinant protein.

NCBI AND UNIPROT INFORMATION

NCBI GI #	21361571
NCBI GeneID	962
NCBI Accession #	NP_001769.2
NCBI GenBank Nucleotide #	NM_001778.3
UniProt Accession #	P09326; Q5U055; Q8MGR0
Molecular Weight	19,429 Da
NCBI Official Full Name	CD48 antigen isoform 1
NCBI Official Synonym Full Names	CD48 molecule
NCBI Official Symbol	CD48
NCBI Official Synonym Symbols	BCM1; BLAST; hCD48; mCD48; BLAST1; SLAMF2; MEM-102
NCBI Protein Information	CD48 antigen
UniProt Protein Name	CD48 antigen
UniProt Gene Name	CD48
UniProt Synonym Gene Names	BCM1; BLAST1; SLAMF2

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/signaling-lymphocytic-activation-molecule-family-antibody-immunohistochemistry-immunocytochemistry-western-blot-elisa/144743>