

Rabbit Xeroderma Pigmentosum, Complementation Group D Polyclonal Antibody | anti-XPD antibody

Xeroderma Pigmentosum, Complementation Group D (XPD) Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA145104	Rabbit	Human, Mouse, Rat	ELISA, Immunohistochemistry, Immunocytochemistry, Western Blot

PRODUCT SYNONYMS

Xeroderma Pigmentosum; Complementation Group D; Polyclonal Antibody; Complementation Group D (XPD) Polyclonal Antibody; CXPB; BTF2 p80; EM9; TTD; ERCC2; TFIIH basal transcription factor complex 80 kDa; De Sanctis-Cacchione; Excision Repair Cross-Complementing Rodent Repair Deficiency 2; anti-XPD antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Specificity	The antibody is a rabbit polyclonal antibody raised against XPD. It has been selected for its ability to recognize XPD in immunohistochemical staining and western blotting.
Purity/Purification	Affinity Chromatography
Form/Format	Supplied as solution form in PBS, pH7.4 containing 0.02% NaN ₃ , 50% glycerol.
Concentration	500ug/mL (varies by lot)
Sequence Length	940
Applicable Applications for anti-XPD antibody	ELISA, IHC (Immunohistochemistry), ICC (Immunocytochemistry), WB (Western Blot)
Immunoglobulin Type	IgG
Conjugate	No Conjugate
Cross Reactivity	Human
Organism Species	Human
Fragment	XPD (Ala404~Leu637)
Quality Control	Content: The quality control contains recombinant XPD (Ala404~Leu637) disposed in loading buffer. Usage: 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate. 5uL per well when used in enhanced chemiluminescent (ECL). Note: The quality control is specifically manufactured as the positive control. Not used for other purposes. Loading Buffer: 100mM Tris(pH8.8), 2% SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, NaN ₃ 0.02%.
Conjugated Antibody	The APC conjugated antibody version of this item is also available as
Preparation and Storage	Store at 4 degree C for frequent use. Stored at -20 degree C to -80 degree C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.

IMAGES



IHC (Immunohiostchemistry)
DAB staining on fromalin fixed paraffin- embedded Kidney tissue



WB (Western Blot)
Western Blot: Sample: Recombinant protein.

NCBI AND UNIPROT INFORMATION

NCBI GI #	15834617
NCBI GeneID	2068
NCBI Accession #	NP_000391.1
NCBI GenBank Nucleotide #	NM_000400.3
UniProt Accession #	P18074; Q2TB78; Q2YDY2; Q7KZU6; Q8N721
Molecular Weight	46,274 Da
NCBI Official Full Name	TFIIH basal transcription factor complex helicase XPD subunit isoform 1
NCBI Official Synonym Full Names	ERCC excision repair 2, TFIIH core complex helicase subunit
NCBI Official Symbol	ERCC2
NCBI Official Synonym Symbols	EM9; TTD; XPD; TTD1; COFS2; TFIIH
NCBI Protein Information	TFIIH basal transcription factor complex helicase XPD subunit
UniProt Protein Name	TFIIH basal transcription factor complex helicase XPD subunit
UniProt Gene Name	ERCC2
UniProt Synonym Gene Names	XPD; XPDC; BTF2 p80; TFIIH 80 kDa subunit; TFIIH p80

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/xeroderma-pigmentosum-complementation-group-d-antibody-immunohistochemistry-immunocytochemistry-western-blot-elisa/145104>