

Rabbit Malic Enzyme 2, NADP+ Dependent Polyclonal Antibody | anti-ME2 antibody

Malic Enzyme 2, NADP+ Dependent, Mitochondrial (ME2) Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA145147	Rabbit	Human, Mouse, Rat	ELISA, Immunohistochemistry, Immunocytochemistry, Western Blot


PRODUCT SYNONYMS

Malic Enzyme 2; NADP+ Dependent; Polyclonal Antibody; Mitochondrial (ME2) Polyclonal Antibody; anti-ME2 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Specificity	The antibody is a rabbit polyclonal antibody raised against ME2. It has been selected for its ability to recognize ME2 in immunohistochemical staining and western blotting.
Purity/Purification	Affinity Chromatography
Form/Format	Supplied as solution form in PBS, pH7.4 containing 0.02% NaN ₃ , 50% glycerol.
Concentration	1mg/ml (varies by lot)
Sequence Length	583
Applicable Applications for anti-ME2 antibody	ELISA, IHC (Immunohistochemistry), ICC (Immunocytochemistry), WB (Western Blot)
Immunoglobulin Type	IgG
Conjugate	No Conjugate
Cross Reactivity	Human
Organism Species	Human
Fragment	ME2 (Gly220~Ala426)
Quality Control	Content: The quality control contains recombinant ME2 (Gly220~Ala426) disposed in loading buffer. Usage: 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate. 5uL per well when used in enhanced chemiluminescent (ECL). Note: The quality control is specifically manufactured as the positive control. Not used for other purposes. Loading Buffer: 100mM Tris(pH8.8), 2% SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, NaN ₃ 0.02%.
Conjugated Antibody	The APC conjugated antibody version of this item is also available as
Preparation and Storage	Store at 4 degree C for frequent use. Stored at -20 degree C to -80 degree C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.


IMAGES




IHC (Immunohistochemistry)
DAB staining on IHC-P; Samples: Human Glioma Tissue




IHC (Immunohistochemistry)
DAB staining on IHC-P; Samples: Human Brain Tissue



IHC (Immunohistochemistry)
DAB staining on IHC-P; Samples: Human Stomach Tissue.



WB (Western Blot)
Western Blot: Sample: Recombinant ME2, Human.



WB (Western Blot)
Western Blot: Sample: Lane1: Human Lung Tissue; Lane2: Human Jurkat Cells; Lane3: Human K562 Cells; Lane4: Human Hela Cells.

NCBI AND UNIPROT INFORMATION

NCBI GI #	270265879
NCBI GeneID	4200
NCBI Accession #	NP_001161807.1
NCBI GenBank Nucleotide #	NM_001168335.1
UniProt Accession #	P23368; Q9BWL6; Q9BYG1; Q9H4B2; B2R8J2
Molecular Weight	53,586 Da

NCBI Official Full Name	NAD-dependent malic enzyme, mitochondrial isoform 2
NCBI Official Synonym Full Names	malic enzyme 2
NCBI Official Symbol	ME2
NCBI Official Synonym Symbols	ODS1
NCBI Protein Information	NAD-dependent malic enzyme, mitochondrial
UniProt Protein Name	NAD-dependent malic enzyme, mitochondrial
UniProt Gene Name	ME2
UniProt Synonym Gene Names	NAD-ME

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/malic-enzyme-2-nadp-dependent-antibody-immunohistochemistry-immunocytochemistry-western-blot-elisa/145147>