

Rabbit Prostaglandin F2 Receptor Negative Regulator Polyclonal Antibody | anti-PTGFRN antibody

Prostaglandin F2 Receptor Negative Regulator (PTGFRN) Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA145159	Rabbit	Human, Mouse, Rat	ELISA, Immunohistochemistry, Immunocytochemistry, Western Blot

PRODUCT SYNONYMS

Prostaglandin F2 Receptor Negative Regulator; Polyclonal Antibody; Prostaglandin F2 Receptor Negative Regulator (PTGFRN) Polyclonal Antibody; CD315; CD9P-1; EWI-F; FPRP; SMAP-6; CD9 partner 1; Glu-Trp-Ile EWI motif-containing protein F; Prostaglandin F2-alpha receptor-associated protein; anti-PTGFRN antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Specificity	The antibody is a rabbit polyclonal antibody raised against PTGFRN. It has been selected for its ability to recognize PTGFRN in immunohistochemical staining and western blotting.
Purity/Purification	Affinity Chromatography
Form/Format	Supplied as solution form in PBS, pH7.4 containing 0.02% NaN3, 50% glycerol.
Concentration	500ug/mL (varies by lot)
Sequence Length	879
Applicable Applications for anti-PTGFRN antibody	ELISA, IHC (Immunohistochemistry), ICC (Immunocytochemistry), WB (Western Blot)
Immunoglobulin Type	IgG
Conjugate	No Conjugate
Cross Reactivity	Human
Organism Species	Human
Fragment	PTGFRN (Pro406~Pro662)
Quality Control	Content: The quality control contains recombinant PTGFRN (Pro406~Pro662) disposed in loading buffer. Usage: 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate. 5uL per well when used in enhanced chemiluminescent (ECL). Note: The quality control is specifically manufactured as the positive control. Not used for other purposes. Loading Buffer: 100mM Tris(pH8.8), 2% SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, NaN3 0.02%.
Conjugated Antibody	The APC conjugated antibody version of this item is also available as
Preparation and Storage	Store at 4 degree C for frequent use. Stored at -20 degree C to -80 degree C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.

IMAGES



IHC (Immunohistochemistry)
DAB staining on formalin fixed paraffin-embedded kidney tissue



WB (Western Blot)
Western Blot: Sample: Recombinant protein.

NCBI AND UNIPROT INFORMATION

NCBI GI #	41152506
NCBI GeneID	5738
NCBI Accession #	NP_065173.2
NCBI GenBank Nucleotide #	NM_020440.3
UniProt Accession #	Q9P2B2; Q5VVU9; Q8N2K6
Molecular Weight	98,556 Da
NCBI Official Full Name	prostaglandin F2 receptor negative regulator
NCBI Official Synonym Full Names	prostaglandin F2 receptor inhibitor
NCBI Official Symbol	PTGFRN
NCBI Official Synonym Symbols	FPRP; CD315; EWI-F; CD9P-1; SMAP-6
NCBI Protein Information	prostaglandin F2 receptor negative regulator
UniProt Protein Name	Prostaglandin F2 receptor negative regulator
UniProt Gene Name	PTGFRN
UniProt Synonym Gene Names	CD9P1; EWIF; FPRP; KIAA1436; CD9P-1; EWI-F

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.